

THE ENRICHED VIETORIS MONAD ON REPRESENTABLE SPACES

DIRK HOFMANN

ABSTRACT. Employing a formal analogy between ordered sets and topological spaces, over the past years we have investigated a notion of cocompleteness for topological, approach and other kind of spaces. In this new context, the down-set monad becomes the filter monad, cocomplete ordered set translates to continuous lattice, distributivity means disconnectedness, and so on. Curiously, the dual(?) notion of completeness does not behave as the mirror image of the one of cocompleteness; and in this paper we have a closer look at complete spaces. In particular, we construct the “up-set monad” on representable spaces (in the sense of L. Nachbin for topological spaces, respectively C. Hermida for multicategories); we show that this monad is of Kock-Zöberlein type; we introduce and study a notion of weighted limit similar to the classical notion for enriched categories; and we describe the Kleisli category of our “up-set monad”. We emphasize that these generic categorical notions and results can be indeed connected to more “classical” topology: for topological spaces, the “up-set monad” becomes the upper Vietoris monad, and the statement “ X is totally cocomplete if and only if X^{op} is totally complete” specialises to O. Wyler’s characterisation of the algebras of the Vietoris monad on compact Hausdorff spaces.

INTRODUCTION

In this paper we continue the work presented in [Hofmann, 2011] on “injective spaces via adjunction” whose fundamental aspect can be described by the slogan *topological spaces are categories*, and therefore can be studied using notions and techniques from (enriched) Category Theory. The use of the word “continue” here is slightly misleading as we do not follow directly the path of [Hofmann, 2011] but rather develop “the second aspect” of this theory. To explain this better, recall that an order relation on a set X defines a monotone map of type

$$X^{\text{op}} \times X \rightarrow 2,$$

and from that one obtains the two Yoneda embeddings

$$X \rightarrow 2^{X^{\text{op}}} =: PX \quad \text{and} \quad X \rightarrow (2^X)^{\text{op}} =: VX.$$

Furthermore, both PX and VX are part of monads $\mathbb{P}$ and $\mathbb{V}$ on Ord (the category of ordered sets and monotone maps) with Eilenberg–Moore categories $\text{Ord}^{\mathbb{P}} \simeq \text{Sup}$ (the category of complete lattices and sup-preserving maps) and $\text{Ord}^{\mathbb{V}} \simeq \text{Inf}$ (the category of complete lattices and inf-preserving maps) respectively. One has full embeddings

$$\text{Ord}_{\mathbb{P}} \rightarrow \text{Ord}^{\mathbb{P}} \quad \text{and} \quad \text{Ord}_{\mathbb{V}} \rightarrow \text{Ord}^{\mathbb{V}}$$

Date: January 28, 2013.

2010 Mathematics Subject Classification. 18A05, 18B10, 18B30, 18B35, 18C15, 18C20, 18D15, 18D20, 54A05, 54A20, 54B20, 54B30.

Key words and phrases. Vietoris construction, enriched category, monad, quantale, topological theory, weighted limit and colimit, adjunction.

Partial financial assistance by FEDER funds through COMPETE – Operational Programme Factors of Competitiveness (Programa Operacional Factores de Competitividade) and by Portuguese funds through the Center for Research and Development in Mathematics and Applications (University of Aveiro) and the Portuguese Foundation for Science and Technology (FCT – Fundação para a Ciência e a Tecnologia), within project PEst-C/MAT/UI4106/2011 with COMPETE number FCOMP-01-0124-FEDER-022690, and the project MONDRIAN under the contract PTDC/EIA-CCO/108302/2008 with COMPETE number FCOMP-01-0124-FEDER-010047 is gratefully acknowledged.

from the Kleisli categories into the Eilenberg–Moore categories, and from that one obtains an equivalence (see [Rosebrugh and Wood, 1994, 2004])

$$\mathrm{kar}(\mathrm{Ord}_{\mathbb{P}}) \simeq \mathrm{CCD}_{\mathrm{sup}} \quad \text{and} \quad \mathrm{kar}(\mathrm{Ord}_{\mathbb{V}}) \simeq \mathrm{CCD}_{\mathrm{inf}}$$

between the idempotent split completion of the Kleisli categories on one side and the categories of completely distributive complete lattice and sup-preserving, respectively inf-preserving, maps on the other. These equivalences restrict to

$$\mathrm{Ord}_{\mathbb{P}} \simeq \mathrm{Tal}_{\mathrm{sup}} \quad \text{and} \quad \mathrm{Ord}_{\mathbb{V}} \simeq \mathrm{Tal}_{\mathrm{inf}},$$

where “Tal” stands for totally algebraic lattice. Finally, having both sides restricted to adjoint morphisms leads to the equivalence

$$\mathrm{Ord}^{\mathrm{op}} \simeq \mathrm{Tal}$$

between the dual category of Ord and the category Tal of totally algebraic lattices and sup- and inf-preserving maps.

In [Hofmann, 2011, 2010] and [Clementino and Hofmann, 2009b] we followed the path on the left described above, but now with geometric objects like topological or approach spaces in lieu of ordered sets (the latter representing “metric” topological spaces, see [Lowen, 1997]). To illustrate this analogy, note that the ultrafilter convergence of a topological space defines a continuous map

$$(UX)^{\mathrm{op}} \times X \rightarrow 2$$

(where 2 is the Sierpiński space and $(UX)^{\mathrm{op}}$ is explained in Section 2). Moreover, the space $(UX)^{\mathrm{op}}$ turns out to be exponentiable, therefore we obtain the Yoneda embedding

$$y_X : X \rightarrow 2^{(UX)^{\mathrm{op}}} =: PX.$$

The “story of cocompleteness” can now be told almost as for ordered sets, and we refer to the above-mentioned papers for detailed information. However, in contrast to the ordered case, the subsequent development of the right side cannot be seen as the dual image of the left side; and it is the aim of this work to explore this path.

The paper is organised as follows.

- In Section 1 we describe our general framework, namely that of a topological theory $\mathcal{T} = (\mathbb{T}, \mathbb{V}, \xi)$ (see [Hofmann, 2007]) consisting of a monad $\mathbb{T} = (T, e, m)$ on $\mathbf{Set}$, a quantale $\mathbb{V}$ and a $\mathbb{T}$ -algebra structure $\xi : T\mathbb{V} \rightarrow \mathbb{V}$ on $\mathbb{V}$. The associated notion of $\mathcal{T}$ -category embodies several types of spaces such as topological, metric or approach spaces, and together with $\mathcal{T}$ -functors and $\mathcal{T}$ -distributors defines the categories $\mathcal{T}\text{-Cat}$ and $\mathcal{T}\text{-Dist}$ respectively. We recall succinctly the main constructions and results, in particular that core-compactness implies tensor-exponentiability. In this context, for topological spaces we give a variation of Alexander’s Subbase Lemma for core-compactness using a simple convergence-theoretic argument (Example 1.9).
- Section 2 is devoted to the important notion of representable $\mathcal{T}$ -categories (Definition 2.5) defined as precisely the pseudo-algebras for a natural lifting of the $\mathbf{Set}$ -monad $\mathbb{T}$ to a monad of Kock–Zöberlein type on $\mathcal{T}\text{-Cat}$. We also introduce the concept of a dualisable $\mathcal{T}$ -graph (Definition 2.11). Our interest in representable $\mathcal{T}$ -categories derives from the fact that these are precisely those $\mathcal{T}$ -categories for which the associated dual $\mathcal{T}$ -graph is a $\mathcal{T}$ -category (Definition 2.13 and Proposition 2.15). For topological T_0 -spaces, the concept of representability specialises to the classical notion of a stably compact space which is closely related to L. Nachbin’s ordered compact Hausdorff spaces.
- In Section 3 we recall the principal facts about weighted colimits and cocomplete $\mathcal{T}$ -categories obtained in [Hofmann, 2011] and [Clementino and Hofmann, 2009b]. We stress that, unlike the classical case of enriched categories, here it is necessary to consider weights with arbitrary codomain, not just the one-element category G . Compared to previous work we change notation and use the designation “totally cocomplete” for a

$\mathcal{T}$ -category admitting all weighed colimits, and say that a $\mathcal{T}$ -category is “cocomplete” whenever it has all those weighed colimits where the codomain of the weight is G .

- From Section 4 on we assume that our monad $\mathbb{T} = (T, e, m)$ satisfies $T1 = 1$. In this section we show that the exponential $\mathbf{V}^X$ is always a dualisable $\mathbb{T}$ -graph and, in the topological case, its dual $(2^X)^{\text{op}}$ turns out to be the upper Vietoris topological space. We point out how this can be used to deduce the classical characterisation of exponentiable topological spaces as precisely the core-compact ones (Example 4.3). The main results of this section state that the construction $X \mapsto (\mathbf{V}^X)^{\text{op}} =: VX$ leads to a monad $\mathbb{V} = (V, \hbar, \omega)$ of Kock-Zöberlein type on both the category $\mathcal{T}\text{-Cat}_{\text{core}}$ of core-compact $\mathcal{T}$ -categories and $\mathcal{T}$ -functors (Theorem 4.23) and $\mathcal{T}\text{-ReprCat}$ of representable $\mathcal{T}$ -categories and pseudo-homomorphisms (Theorem 4.13).
- In Section 5 we analyse the notion of weighted limit in $\mathcal{T}$ -categories.
- Section 6 lifts the classical adjunction between “up-sets” and “down-sets” (see [Wood, 2004, Section 5]) into the realm of $\mathcal{T}$ -categories.
- In Section 7 we introduce totally complete $\mathcal{T}$ -categories and show that they are precisely the duals of totally cocomplete $\mathcal{T}$ -categories (Theorem 7.4 and Examples 7.6).
- Finally, in Section 8 we give a characterisation of the morphisms of the Kleisli category $\mathcal{T}\text{-ReprCat}_{\mathbb{V}}$ of $\mathbb{V}$. We also observe how the notion of an Esakia space arises naturally in this context via splitting idempotents of the full subcategory of $\mathcal{T}\text{-ReprCat}_{\mathbb{V}}$ defined by all $\mathbb{T}$ -algebras. We find it worthwhile to mention that this implies in particular that the category $\mathbf{Heyt}_{\perp, \mathbb{V}}$ of Heyting algebras and finite suprema preserving maps is the idempotent split completion of the category $\mathbf{Bool}_{\perp, \mathbb{V}}$ of Boolean algebras and finite suprema preserving maps (Example 8.11).

1. THE SETTING

In this paper we will work with $\mathcal{T}$ -categories, $\mathcal{T}$ -functors and $\mathcal{T}$ -distributors, for a (strict) topological theory $\mathcal{T}$. Below we recall some of the main facts and refer to [Hofmann, 2007], [Hofmann, 2011] and [Clementino and Hofmann, 2009b] for details.

Definition 1.1. A topological theory $\mathcal{T} = (\mathbb{T}, \mathbf{V}, \xi)$ consists of:

- (1) a monad $\mathbb{T} = (T, e, m)$ on $\mathbf{Set}$ (with multiplication m and unit e),
- (2) a commutative and unital quantale $\mathbf{V} = (\mathbf{V}, \otimes, k)$,
- (3) a function $\xi : T\mathbf{V} \rightarrow \mathbf{V}$,

such that

- (a) T preserves weak pullbacks and each naturality square of m is a weak pullback,
- (b) the pair $(\mathbf{V}, \xi)$ is an Eilenberg–Moore algebra for $\mathbb{T}$ and the monoid structure on $\mathbf{V}$ in $(\mathbf{Set}, \times, 1)$ lifts to a monoid structure on $(\mathbf{V}, \xi)$ in $(\mathbf{Set}^{\mathbb{T}}, \times, 1)$, that is, the following diagrams have to commute:

$$\begin{array}{ccccc}
 T(\mathbf{V} \times \mathbf{V}) & \xrightarrow{T(-\otimes-)} & T\mathbf{V} & \xleftarrow{T(k)} & T1 \\
 \downarrow \langle \xi \cdot T\pi_1, \xi \cdot T\pi_2 \rangle & & \downarrow \xi & & \downarrow ! \\
 \mathbf{V} \times \mathbf{V} & \xrightarrow{-\otimes-} & \mathbf{V} & \xleftarrow{k} & 1,
 \end{array}$$

- (c) writing $P_{\mathbf{V}} : \mathbf{Set} \rightarrow \mathbf{Ord}$ for the functor that sends a function $f : X \rightarrow Y$ to the left adjoint of the “inverse image” function $f^{-1} : \mathbf{V}^Y \rightarrow \mathbf{V}^X$, $\varphi \mapsto \varphi \cdot f$ (where $\mathbf{V}^X$ is the set of functions from X to $\mathbf{V}$, with pointwise order), the functions $\xi_X : \mathbf{V}^X \rightarrow \mathbf{V}^{TX}$, $f \mapsto \xi \cdot Tf$ (for X in $\mathbf{Set}$) are the components of a natural transformation $(\xi_X)_X : P_{\mathbf{V}} \rightarrow P_{\mathbf{V}}T$.

Remark 1.2. As shown in [Hofmann, 2007, Lemma 3.2], the *internal hom* in $\mathbf{V}$ defined by

$$x \otimes y \leq z \iff x \leq \text{hom}(y, z)$$

automatically satisfies

$$\begin{array}{ccc}
 T(\mathbf{V} \times \mathbf{V}) & \xrightarrow{T(\text{hom})} & T\mathbf{V} \\
 \langle \xi \cdot T\pi_1, \xi \cdot T\pi_2 \rangle \downarrow & \geq & \downarrow \xi \\
 \mathbf{V} \times \mathbf{V} & \xrightarrow{\text{hom}} & \mathbf{V}.
 \end{array}$$

Remark 1.3. Our notation differs here from [Hofmann, 2007] where ξ is only assumed to be a lax Eilenberg–Moore structure on $\mathbf{V}$ and the diagrams in (b) are only required to commute laxly. A theory satisfying the stronger conditions above is called *strict topological theory* there. However, in this paper all theories are assumed to be strict, therefore we simply use the term “topological theory”.

Throughout this paper we will assume that a topological theory $\mathcal{T} = (\mathbb{T}, \mathbf{V}, \xi)$ is given. Moreover, we will *always assume that $\mathbf{V}$ is non-trivial*, that is, $\perp \neq k$. Consequently, since $\mathbf{V}$ is assumed to be a $\mathbb{T}$ -algebra, the monad $\mathbb{T}$ must be non-trivial. We recall here that there are two trivial monads $\mathbb{T} = (T, e, m)$ on **Set**: one with $TX = 1$ for every set X , and one with $TX = 1$ for every non-empty set and $T\emptyset = \emptyset$. A monad $\mathbb{T} = (T, e, m)$ different from these two is called non-trivial. For a non-trivial monad, T is faithful and e is point-wise injective (see [Manes, 1976], for instance).

Our leading examples are the following:

Examples 1.4.

- (1) For any quantale $\mathbf{V}$ we can consider the theory whose monad-part is the identity monad on **Set** and where $\xi : \mathbf{V} \rightarrow \mathbf{V}$ is the identity function. We write this trivial topological theory as $\mathcal{J}_{\mathbf{V}}$ or simply as $\mathcal{J}$.
- (2) Let $\mathbf{V}$ be the 2-element chain 2 , and consider the ultrafilter monad $\mathbb{U} = (U, e, m)$ on **Set**. This together with the “identity” function $\xi : U2 \rightarrow 2$ is a topological theory which we denote by $\mathcal{U}_2$.
- (3) More general, for a non-trivial monad $\mathbb{T} = (T, e, m)$ on **Set** where T preserves weak pullbacks and each naturality square of m is a weak pullback and every completely distributive complete lattice $\mathbf{V}$ (considering $\otimes = \wedge$ and $k = \top$), $(\mathbb{T}, \mathbf{V}, \xi)$ is a topological theory where

$$\xi : T\mathbf{V} \rightarrow \mathbf{V}, \quad \mathfrak{x} \mapsto \bigvee \{v \in \mathbf{V} \mid \mathfrak{x} \in T(\uparrow v)\}.$$

- (4) In particular, for the ultrafilter monad $\mathbb{U} = (U, e, m)$ on **Set** and the complete lattice $[0, \infty]$ ordered by the “greater or equal” relation $\geq$ (so that the infimum of two numbers is their maximum and the supremum of $S \subseteq [0, \infty]$ is given by $\inf S$), we write $\mathbf{P}_{\wedge} = ([0, \infty], \max, 0)$ for the corresponding quantale and $\mathcal{U}_{\mathbf{P}_{\wedge}} = (\mathbb{U}, \mathbf{P}_{\wedge}, \xi)$ for the corresponding theory where

$$\xi : U([0, \infty]) \rightarrow [0, \infty], \quad \mathfrak{x} \mapsto \inf \{v \in [0, \infty] \mid [0, v] \in \mathfrak{x}\}.$$

Also note that

$$\text{hom}(u, v) = \begin{cases} 0 & \text{if } u \geq v, \\ v & \text{otherwise} \end{cases}$$

in $\mathbf{P}_{\wedge}$.

- (5) Let $\mathbf{V}$ be the quantale $\mathbf{P}_{+} = ([0, \infty], +, 0)$ of extended non-negative real numbers ordered by the “greater or equal” relation (see [Lawvere, 1973]), and consider again the ultrafilter monad $\mathbb{U} = (U, e, m)$ on **Set**. Together with the function $\xi : U([0, \infty]) \rightarrow [0, \infty]$ as above this makes up a topological theory, denoted as $\mathcal{U}_{\mathbf{P}_{+}}$. For later use we record here that the internal hom of the quantale $\mathbf{P}_{+}$ is given by truncated minus:

$$\text{hom}(u, v) = v \ominus u := \max\{v - u, 0\}.$$

(6) For any quantale $\mathbf{V}$, the word monad $\mathbb{L} = (L, e, m)$ on $\mathbf{Set}$ together with the function

$$\xi : L(\mathbf{V}) \rightarrow \mathbf{V}, (v_1, \dots, v_n) \mapsto v_1 \otimes \dots \otimes v_n, () \mapsto k$$

determine a topological theory.

Since some of our principal examples involve the ultrafilter monad, we note here two important results.

Proposition 1.5. *Let X be a set, $\mathfrak{f}$ a filter and $\mathfrak{j}$ an ideal on X with $\mathfrak{f} \cap \mathfrak{j} = \emptyset$. Then there exists an ultrafilter $\mathfrak{x}$ on X with $\mathfrak{f} \subseteq \mathfrak{x}$ and $\mathfrak{x} \cap \mathfrak{j} = \emptyset$.*

Proof. See [Stone, 1938, Theorem 6], for instance. $\square$

Theorem 1.6 ([Manes, 1969]). *The Eilenberg–Moore category $\mathbf{Set}^{\mathbb{U}}$ of the ultrafilter monad on $\mathbf{Set}$ is equivalent to the category $\mathbf{CompHaus}$ of compact Hausdorff spaces and continuous maps.*

Every topological theory allows for a number of constructions and definitions which were succinctly recalled in [Clementino and Hofmann, 2009b]. Below we give a slightly revised version of [Clementino and Hofmann, 2009b, Section 1].

I. The quantaloid $\mathbf{V}\text{-Rel}$ (see [Betti *et al.*, 1983]) has sets as objects, and a morphism $r : X \multimap Y$ from X to Y is a $\mathbf{V}$ -relation $r : X \times Y \rightarrow \mathbf{V}$ (also called $\mathbf{V}$ -matrix). The composition of $\mathbf{V}$ -relations $r : X \multimap Y$ and $s : Y \multimap Z$ is defined as matrix multiplication

$$s \cdot r(x, z) = \bigvee_{y \in Y} r(x, y) \otimes s(y, z),$$

and the identity arrow $1_X : X \multimap X$ is the $\mathbf{V}$ -relation which sends all diagonal elements (x, x) to k and all other elements to the bottom element $\perp$ of $\mathbf{V}$. The set $\mathbf{V}\text{-Rel}(X, Y)$ of all $\mathbf{V}$ -relations from X to Y becomes a complete ordered set by putting

$$r \leq r' \text{ whenever } \forall x \in X \forall y \in Y. r(x, y) \leq r'(x, y),$$

for $\mathbf{V}$ -relations $r, r' : X \multimap Y$; composition from either side preserves this order.

The category $\mathbf{V}\text{-Rel}$ has an involution $(r : X \multimap Y) \mapsto (r^\circ : Y \multimap X)$ where $r^\circ(y, x) = r(x, y)$, satisfying

$$1_X^\circ = 1_X, \quad (s \cdot r)^\circ = r^\circ \cdot s^\circ, \quad r^{\circ\circ} = r,$$

as well as $r^\circ \leq s^\circ$ whenever $r \leq s$. Furthermore, there is a faithful functor

$$\mathbf{Set} \rightarrow \mathbf{V}\text{-Rel}, (f : X \rightarrow Y) \mapsto (f : X \multimap Y)$$

sending a map $f : X \rightarrow Y$ to its graph $f : X \multimap Y$ defined by

$$f(x, y) = \begin{cases} k & \text{if } f(x) = y, \\ \perp & \text{else.} \end{cases}$$

In the sequel we will not distinguish between the function f and the $\mathbf{V}$ -relation f and simply write $f : X \rightarrow Y$. We also note that $f \dashv f^\circ$ in the quantaloid $\mathbf{V}\text{-Rel}$.

Let $t : X \multimap Z$ be a $\mathbf{V}$ -relation. The composition functions

$$- \cdot t : \mathbf{V}\text{-Rel}(Z, Y) \rightarrow \mathbf{V}\text{-Rel}(X, Y) \quad \text{and} \quad t \cdot - : \mathbf{V}\text{-Rel}(Y, X) \rightarrow \mathbf{V}\text{-Rel}(Y, Z).$$

preserve suprema and therefore have respective right adjoints

$$(-) \bullet - t : \mathbf{V}\text{-Rel}(X, Y) \rightarrow \mathbf{V}\text{-Rel}(Z, Y) \quad \text{and} \quad t \bullet (-) : \mathbf{V}\text{-Rel}(Y, Z) \rightarrow \mathbf{V}\text{-Rel}(Y, X).$$

Here, for $\mathbf{V}$ -relations $r : X \multimap Y$ and $s : Y \multimap Z$,

$$(r \bullet - t)(z, y) = \bigwedge_{x \in X} \text{hom}(t(x, z), r(x, y)) \quad (t \bullet s)(y, x) = \bigwedge_{z \in Z} \text{hom}(t(x, z), s(y, z)).$$

We call $r \bullet - t$ the *extension of r along t* , and $t \bullet s$ the *lifting of r along t* .

II. The **Set**-functor T extends to a 2-functor $T_\xi : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$. To each $\mathbf{V}$ -relation $r : X \times Y \rightarrow \mathbf{V}$, T_ξ assigns the $\mathbf{V}$ -relation $T_\xi r : TX \times TY \rightarrow \mathbf{V}$ such that, for every map $s : TX \times TY \rightarrow \mathbf{V}$,

$$\xi \cdot Tr \leq s \cdot \langle T\pi_1, T\pi_2 \rangle \Leftrightarrow T_\xi r \leq s :$$

$$\begin{array}{ccc} & TX \times TY & \\ \langle T\pi_1, T\pi_2 \rangle \uparrow & \searrow T_\xi r & \\ T(X \times Y) & \xrightarrow{\xi \cdot Tr} & \mathbf{V} \end{array} \quad \leq$$

In other words, regarding TX , TY and $TX \times TY$ as discrete ordered sets, $T_\xi r$ is the left Kan extension in **Ord** of $\xi \cdot Tr$ along $\langle T\pi_1, T\pi_2 \rangle$. Hence, for $\mathfrak{x} \in TX$ and $\mathfrak{y} \in TY$,

$$T_\xi r(\mathfrak{x}, \mathfrak{y}) = \bigvee \left\{ \xi \cdot Tr(\mathfrak{w}) \mid \mathfrak{w} \in T(X \times Y), T\pi_1(\mathfrak{w}) = \mathfrak{x}, T\pi_2(\mathfrak{w}) = \mathfrak{y} \right\}.$$

The 2-functor T_ξ preserves the involution in the sense that $T_\xi(r^\circ) = T_\xi(r)^\circ$ (and we write $T_\xi r^\circ$) for each $\mathbf{V}$ -relation $r : X \multimap Y$, m becomes a natural transformation $m : T_\xi T_\xi \rightarrow T_\xi$ and e an op-lax natural transformation $e : 1 \rightarrow T_\xi$, that is, $e_Y \cdot r \leq T_\xi r \cdot e_X$ for all $r : X \multimap Y$ in $\mathbf{V}\text{-Rel}$.

For $\mathcal{T} = \mathcal{U}_2$, the extension above coincides with the one given in [Barr, 1970]; and for $\mathcal{T} = \mathcal{U}_{\mathbf{P}_+}$ and $\mathcal{T} = \mathcal{U}_{\mathbf{P}_\wedge}$ one obtains

$$U_\xi r(\mathfrak{x}, \mathfrak{y}) = \sup_{A \in \mathfrak{x}, B \in \mathfrak{y}} \inf_{x \in A, y \in B} r(x, y)$$

for all $r : X \multimap Y$, $\mathfrak{x} \in UX$ and $\mathfrak{y} \in UY$ (see also [Clementino and Tholen, 2003]).

Different methods for extending **Set**-functors to **Rel** can be found in [Seal, 2005; Schubert and Seal, 2008; Seal, 2009].

III. $\mathbf{V}$ -relations of the form $\alpha : TX \multimap Y$, called $\mathcal{T}$ -relations and denoted by $\alpha : X \multimap Y$, will play an important role here. Given two $\mathcal{T}$ -relations $\alpha : X \multimap Y$ and $\beta : Y \multimap Z$, their *Kleisli convolution* $\beta \circ \alpha : X \multimap Z$ is defined as

$$\beta \circ \alpha = \beta \cdot T_\xi \alpha \cdot m_X^\circ.$$

This operation is associative and has the $\mathcal{T}$ -relation $e_X^\circ : X \multimap X$ as a lax identity:

$$a \circ e_X^\circ = a \quad \text{and} \quad e_Y^\circ \circ a \geq a,$$

for any $a : X \multimap Y$.

IV. Those $\mathcal{T}$ -relations satisfying the usual unit and composition axioms of a category define $\mathcal{T}$ -categories: a $\mathcal{T}$ -category is a pair (X, a) consisting of a set X and a $\mathcal{T}$ -relation $a : X \multimap X$ on X such that

$$e_X^\circ \leq a \quad \text{and} \quad a \circ a \leq a.$$

Expressed elementwise, these conditions become

$$k \leq a(e_X(x), x) \quad \text{and} \quad T_\xi a(\mathfrak{X}, \mathfrak{x}) \otimes a(\mathfrak{x}, x) \leq a(m_X(\mathfrak{X}), x)$$

for all $\mathfrak{X} \in TTX$, $\mathfrak{x} \in TX$ and $x \in X$. We refer to the first condition as *reflexivity* and to the second one as *transitivity*. A function $f : X \rightarrow Y$ between $\mathcal{T}$ -categories (X, a) and (Y, b) is a $\mathcal{T}$ -functor if $f \cdot a \leq b \cdot Tf$, which in pointwise notation reads as

$$a(\mathfrak{x}, x) \leq b(Tf(\mathfrak{x}), f(x))$$

for all $\mathfrak{x} \in TX$, $x \in X$. The category of $\mathcal{T}$ -categories and $\mathcal{T}$ -functors is denoted by

$$\mathcal{T}\text{-Cat}.$$

If $\mathcal{T} = \mathcal{J}_V$ is an identity theory, for a quantale V , then a $\mathcal{T}$ -category is just a V -category and $\mathcal{T}$ -functor means V -functor (in the sense of [Eilenberg and Kelly, 1966]). Therefore we write V -category instead of $\mathcal{J}_V$ -category, V -functor instead of $\mathcal{J}_V$ -functor, and

V -Cat

instead of $\mathcal{J}_V$ -Cat. We also recall that $2\text{-Cat} \simeq \text{Ord}$, $P_+\text{-Cat} \simeq \text{Met}$ (the category of generalised metric spaces and non-expansive maps, see [Lawvere, 1973]) and $P_\wedge\text{-Cat} \simeq \text{UMet}$ (the category of generalised ultrametric spaces and non-expansive maps). Our principal examples are the ultrafilter theories $\mathcal{U}_2$ and $\mathcal{U}_{P_+}$: the main result of [Barr, 1970] states that $\mathcal{U}_2\text{-Cat}$ is isomorphic to the category **Top** of topological spaces and continuous maps, and in [Clementino and Hofmann, 2003] it is shown that $\mathcal{U}_{P_+}\text{-Cat}$ is isomorphic to the category **App** of approach spaces and non-expansive maps [Lowen, 1989] (regarding notation and results about approach spaces we refer to [Lowen, 1997]). The category $\mathcal{U}_{P_\wedge}\text{-Cat}$ can be identified with the full subcategory **UApp** of **App** defined by all those approach spaces (X, a) which satisfy

$$\max(U_\xi a(\mathfrak{X}, \mathfrak{x}), a(\mathfrak{x}, x)) \geq a(m_X(\mathfrak{X}), x),$$

for all $\mathfrak{X} \in UUX$, $\mathfrak{x} \in UX$ and $x \in X$. In the sequel we always refer to these presentations when talking about **Ord**, **Met**, **UMet**, **Top**, **App** or **UApp**.

V. The forgetful functor $\mathcal{T}\text{-Cat} \rightarrow \text{Set}$, $(X, a) \mapsto X$ is topological, hence it has a left and a right adjoint. In particular, the free $\mathcal{T}$ -category on a set X is given by (X, e_X°) . The $\mathcal{T}$ -category $G = (1, e_1^\circ)$ is a generator in $\mathcal{T}\text{-Cat}$. Furthermore, there is a canonical forgetful functor $\mathcal{T}\text{-Cat} \rightarrow V\text{-Cat}$, commuting with the forgetful functors to **Set**, which sends a $\mathcal{T}$ -category (X, a) to the V -category (X, a_0) where $a_0 = a \cdot e_X$; and $\mathcal{T}\text{-Cat} \rightarrow V\text{-Cat}$ has a concrete left adjoint which sends a V -category (X, c) to $(X, e_X^\circ \cdot T_\xi c)$.

VI. A $\mathcal{T}$ -relation $\varphi : X \multimap Y$ between $\mathcal{T}$ -categories $X = (X, a)$ and $Y = (Y, b)$ is a $\mathcal{T}$ -distributor, denoted as $\varphi : X \multimap Y$, if $\varphi \circ a \leq \varphi$ and $b \circ \varphi \leq \varphi$. Note that we always have $\varphi \circ a \geq \varphi$ and $b \circ \varphi \geq \varphi$, so that the $\mathcal{T}$ -distributor conditions above are in fact equalities. $\mathcal{T}$ -categories and $\mathcal{T}$ -distributors form a 2-category, denoted by

$\mathcal{T}\text{-Dist}$,

with Kleisli convolution as composition and with the 2-categorical structure inherited from **V-Rel**. The identity in $\mathcal{T}\text{-Dist}$ on a $\mathcal{T}$ -category $X = (X, a)$ is given by $a : X \multimap X$. As before, we write

$V\text{-Dist}$

whenever $\mathcal{T} = \mathcal{J}_V$ is an identity theory, and use $\varphi : X \multimap Y$ instead of $\varphi : X \multimap Y$ in this case.

VII. Each $\mathcal{T}$ -functor $f : (X, a) \rightarrow (Y, b)$ induces an adjunction

$$f_\circ \dashv f^\circ$$

in $\mathcal{T}\text{-Dist}$, with $f_\circ : X \multimap Y$ and $f^\circ : Y \multimap X$ defined as $f_\circ = b \cdot T f$ and $f^\circ = f^\circ \cdot b$ respectively. In fact, these assignments define functors

$$(-)_\circ : \mathcal{T}\text{-Cat} \rightarrow \mathcal{T}\text{-Dist} \quad \text{and} \quad (-)^\circ : \mathcal{T}\text{-Cat}^{\text{op}} \rightarrow \mathcal{T}\text{-Dist},$$

where $X_\circ = X = X^\circ$. More generally, the definition of $f_\circ$ and of f° makes sense for any map $f : X \rightarrow Y$ between $\mathcal{T}$ -categories, not just for $\mathcal{T}$ -functors; but then $f_\circ : X \multimap Y$ and $f^\circ : Y \multimap X$ are in general only $\mathcal{T}$ -relations. However:

$$f \text{ is a } \mathcal{T}\text{-functor} \iff f_\circ \text{ is a } \mathcal{T}\text{-distributor} \iff f^\circ \text{ is a } \mathcal{T}\text{-distributor}. \quad (1)$$

A $\mathcal{T}$ -functor $f : (X, a) \rightarrow (Y, b)$ is called *fully faithful* if $f^\circ \circ f_\circ = 1_X^\circ$. Note that f is fully faithful if and only if, for all $\mathfrak{x} \in TX$ and $x \in X$, $a(\mathfrak{x}, x) = b(Tf(\mathfrak{x}), f(x))$.

For a V -functor $f : X \rightarrow Y$ we will, however, use the traditional notation $f_* : X \multimap Y$ and $f^* : Y \multimap X$. This distinction is convenient since at some occasions we will consider simultaneously

the $\mathcal{T}$ -distributor $f_{\otimes} : (X, a) \multimap (Y, b)$ (induced by the $\mathcal{T}$ -functor $f : (X, a) \rightarrow (Y, b)$) and the $\mathbf{V}$ -distributor $f_* : (X, a_0) \multimap (Y, b_0)$ (induced by the underlying $\mathbf{V}$ -functor $f : (X, a_0) \rightarrow (Y, b_0)$).

The category $\mathcal{T}\text{-Cat}$ becomes a 2-category by transporting the order-structure on hom-sets from $\mathcal{T}\text{-Dist}$ to $\mathcal{T}\text{-Cat}$ via the functor $(-)^{\otimes} : \mathcal{T}\text{-Cat}^{\text{op}} \rightarrow \mathcal{T}\text{-Dist}$: for $\mathcal{T}$ -functors $f, g : (X, a) \rightarrow (Y, b)$ we define (see [Hofmann and Tholen, 2010, Lemma 4.7])

$$\begin{aligned} f \leq g \text{ in } \mathcal{T}\text{-Cat} &: \iff f^{\otimes} \leq g^{\otimes} \text{ in } \mathcal{T}\text{-Dist} && \iff g_{\otimes} \leq f_{\otimes} \text{ in } \mathcal{T}\text{-Dist} \\ &\iff \forall x \in X. k \leq b_0(f(x), g(x)) \\ &\iff f^* \leq g^* \text{ in } \mathbf{V}\text{-Dist} && \iff g_* \leq f_* \text{ in } \mathbf{V}\text{-Dist}. \end{aligned}$$

We call $f, g : X \rightarrow Y$ *equivalent*, and write $f \simeq g$, if $f \leq g$ and $g \leq f$. Hence, $f \simeq g$ if and only if $f^{\otimes} = g^{\otimes}$ if and only if $f_{\otimes} = g_{\otimes}$. A $\mathcal{T}$ -category X is called *separated* (see [Hofmann and Tholen, 2010] for details) whenever $f \simeq g$ implies $f = g$, for all $\mathcal{T}$ -functors $f, g : Y \rightarrow X$ with codomain X . One easily verifies that it is enough to consider the case $Y = G$, so that X is separated if and only if the ordered set $\mathcal{T}\text{-Cat}(G, X)$ is anti-symmetric. The full subcategory of $\mathcal{T}\text{-Cat}$ consisting of all separated $\mathcal{T}$ -categories is denoted by

$$\mathcal{T}\text{-Cat}_{\text{sep}}.$$

The 2-categorical structure on $\mathcal{T}\text{-Cat}$ allows us to consider adjoint $\mathcal{T}$ -functors: a $\mathcal{T}$ -functor $f : X \rightarrow Y$ is *left adjoint* if there exists a $\mathcal{T}$ -functor $g : Y \rightarrow X$ such that $1_X \leq g \cdot f$ and $1_Y \geq f \cdot g$. Considering the corresponding $\mathcal{T}$ -distributors, f is left adjoint to g in $\mathcal{T}\text{-Cat}$ if and only if $g_{\otimes} \dashv f_{\otimes}$ in $\mathcal{T}\text{-Dist}$, that is, if and only if $f_{\otimes} = g^{\otimes}$.

VIII. For a $\mathcal{T}$ -distributor $\alpha : X \multimap Y$, the composition function $- \circ \alpha$ has a right adjoint

$$(-) \circ \alpha \dashv (-) \multimap \alpha$$

where, for a given $\mathcal{T}$ -distributor $\gamma : X \multimap Z$, the *extension* $\gamma \multimap \alpha : Y \multimap Z$ is constructed in $\mathbf{V}\text{-Rel}$ as the extension $\gamma \multimap \alpha = \gamma \bullet (T_{\xi}\alpha \cdot m_X^{\circ})$.

$$\begin{array}{ccc} TX & \xrightarrow{\gamma} & Z \\ m_X^{\circ} \downarrow & \nearrow & \\ TT X & & \\ T_{\xi}\alpha \downarrow & & \\ TY & & \end{array}$$

Unfortunately, *in general liftings need not exist* in $\mathcal{T}\text{-Dist}$ (see [Hofmann and Stubbe, 2011, Example 1.7]).

IX. The tensor product on $\mathbf{V}$ can be transported to $\mathcal{T}\text{-Cat}$ by putting

$$(X, a) \otimes (Y, b) = (X \times Y, c),$$

with

$$c(\mathfrak{w}, (x, y)) = a(T\pi_1(\mathfrak{w}), x) \otimes b(T\pi_2(\mathfrak{w}), y),$$

where $\mathfrak{w} \in T(X \times Y)$, $x \in X$, $y \in Y$. The $\mathcal{T}$ -category $E = (1, k)$ is a $\otimes$ -neutral object, where 1 is a singleton set and $k : T1 \times 1 \rightarrow \mathbf{V}$ the constant relation with value $k \in \mathbf{V}$. In general, this construction does not result in a closed structure on $\mathcal{T}\text{-Cat}$; however, it does so when defined in the larger category $\mathcal{T}\text{-Gph}$ of $\mathcal{T}$ -graphs and $\mathcal{T}$ -graph morphisms. Here a $\mathcal{T}$ -graph (see [Clementino et al., 2003]) is a pair (X, a) consisting of a set X and a $\mathbf{V}$ -relation $a : TX \times X \rightarrow \mathbf{V}$ which is only required to satisfy

$$k \leq a(e_X(x), x),$$

for all $x \in X$; $\mathcal{T}$ -graph morphisms are defined as $\mathcal{T}$ -functors. There is an obvious full embedding

$$\mathcal{T}\text{-Cat} \hookrightarrow \mathcal{T}\text{-Gph}.$$

For $\mathcal{T}$ -graphs $X = (X, a)$ and $Y = (Y, b)$, $X \otimes Y$ is defined as above, but now $X \otimes - : \mathcal{T}\text{-Gph} \rightarrow \mathcal{T}\text{-Gph}$ has a right adjoint $(-)^X : \mathcal{T}\text{-Gph} \rightarrow \mathcal{T}\text{-Gph}$ (see [Hofmann, 2007]) where the structure d on

$$Y^X = \{f : X \rightarrow Y \mid f \text{ is a } \mathcal{T}\text{-functor of type } G \otimes X \rightarrow Y\}$$

is given by

$$d(\mathfrak{p}, h) = \bigwedge_{\substack{\mathfrak{q} \in T(Y^X \times X), x \in X \\ \mathfrak{q} \mapsto \mathfrak{p}}} \text{hom}(a(T\pi_2(\mathfrak{q}), x), b(\text{TeV}(\mathfrak{q}), h(x))).$$

Here ev denotes the evaluation map $\text{ev} : Y^X \times X \rightarrow Y, (h, x) \mapsto h(x)$. The following result can be found in [Hofmann, 2007].

Proposition 1.7. *Let $X = (X, a)$ be a $\mathcal{T}$ -category with $a \cdot T_\xi a = a \cdot m_X$. Then, for each $\mathcal{T}$ -category Y , the structure d on Y^X is transitive. Hence, $X \otimes - : \mathcal{T}\text{-Cat} \rightarrow \mathcal{T}\text{-Cat}$ has a right adjoint $(-)^X : \mathcal{T}\text{-Cat} \rightarrow \mathcal{T}\text{-Cat}$. Moreover, for $h, h' \in Y^X$,*

$$d(e_{Y^X}(h'), h) = \bigwedge_{x \in X} b(e_Y(h'(x)), h(x)).$$

Definition 1.8. A $\mathcal{T}$ -category $X = (X, a)$ is called *core-compact* whenever $a \cdot T_\xi a = a \cdot m_X$.

Example 1.9. The designation “core-compact” is motivated by the case of topological spaces. Classically, a topological space X with topology $\mathcal{O}$ is called core-compact whenever $x \in U \in \mathcal{O}$ implies that there exists some $V \in \mathcal{O}$ with $x \in V$ and V is *relatively compact* in U ; the latter meaning that very open cover of U contains a finite sub-cover of V , or, equivalently, every ultrafilter on V has a convergence point in U . It is shown in [Pisani, 1999] that X is core-compact if and only if its convergence structure $a : UX \rightarrow X$ satisfies $a \cdot U_\xi a = a \cdot m_X$. We find it worthwhile to note that the proof of the implication “core-compact $\Rightarrow a \cdot U_\xi a = a \cdot m_X$ ” can be adapted to subbases, under a certain condition. More in detail, for a set X equipped with a subset $\mathcal{B}$ of the powerset of X (no axioms required), if $(X, \mathcal{B})$ is core-compact (defined as for topological spaces), then the induced convergence $a : UX \rightarrow X$ (defined as for topological spaces) satisfies $a \cdot U_\xi a = a \cdot m_X$ provided that every ultrafilter has a smallest convergence point with respect to the convergence a and the order relation $a \cdot e_X$. Since the topology induced by $\mathcal{B}$ has the same convergence as $\mathcal{B}$, one obtains a variation of Alexander’s Sub-Base Lemma: A topological space where every ultrafilter has a smallest convergence point is core-compact if it is core-compact with respect to a sub-basis. We will apply this principle in Example 4.3. We also note that this argument works for any property of topological spaces which can be equivalently expressed in terms of opens and in terms of ultrafilter convergence, without using the axioms of a topology; in particular in the classical case of compactness.

2. REPRESENTABLE $\mathcal{T}$ -CATEGORIES AND DUALISATION

In [Clementino and Hofmann, 2009a] we introduced a notion of dual $\mathcal{T}$ -category as a crucial step towards the Yoneda lemma and related results. The basic idea is to associate to a $\mathcal{T}$ -category X a $\mathbf{V}$ -category MX which still contains all information about the $\mathcal{T}$ -categorical structure of X , and then use the usual dualisation of $\mathbf{V}$ -categories. Later, in [Hofmann, 2010; Gutierrez and Hofmann, 2012], we noted already that this construction is closely related to Nachbin’s ordered compact Hausdorff spaces [Nachbin, 1950] as presented in [Tholen, 2009]. In this section we continue this path and introduce a class of $\mathcal{T}$ -categories (designated as representable $\mathcal{T}$ -categories) which naturally admit a dual.

Recall from [Tholen, 2009] that the **Set**-monad $\mathbb{T} = (T, e, m)$ admits a natural extension to a monad on $\mathbf{V}\text{-Cat}$, in the sequel also denoted as $\mathbb{T} = (T, e, m)$. Here the functor $T : \mathbf{V}\text{-Cat} \rightarrow \mathbf{V}\text{-Cat}$ sends a $\mathbf{V}$ -category (X, a_0) to $(TX, T_\xi a_0)$, and with this definition $e_X : X \rightarrow TX$ and $m_X : TTX \rightarrow TX$ become $\mathbf{V}$ -functors for each $\mathbf{V}$ -category X . We also note that $T : \mathbf{V}\text{-Cat} \rightarrow \mathbf{V}\text{-Cat}$ is actually a 2-functor: if $f^* \leq g^*$, then $(Tf)^* = T_\xi(f^*) \leq T_\xi(g^*) = (Tg)^*$.

Eilenberg–Moore algebras for this monad can be described as triples (X, a_0, α) where (X, a_0) is a $\mathbf{V}$ -category and (X, α) is an algebra for the $\mathbf{Set}$ -monad $\mathbb{T}$ such that $\alpha : T(X, a_0) \rightarrow (X, a_0)$ is a $\mathbf{V}$ -functor. For $\mathbb{T}$ -algebras (X, a_0, α) and (Y, b_0, β) , a map $f : X \rightarrow Y$ is a homomorphism $f : (X, a_0, \alpha) \rightarrow (Y, b_0, \beta)$ precisely if f preserves both structures, that is, whenever $f : (X, a_0) \rightarrow (Y, b_0)$ is a $\mathbf{V}$ -functor and $f : (X, \alpha) \rightarrow (Y, \beta)$ is a $\mathbb{T}$ -homomorphism. Since the extension T_ξ of T commutes with the involution $(-)^{\circ}$, with (X, a_0, α) also (X, a_0°, α) is a $\mathbb{T}$ -algebra.

Example 2.1. It follows from Remark 1.2 that, for every topological theory $\mathcal{T} = (\mathbb{T}, \mathbf{V}, \xi)$, the internal hom in $\mathbf{V}$ combined with the $\mathbb{T}$ -algebra structure ξ induces the Eilenberg–Moore algebra $\mathbf{V} = (\mathbf{V}, \text{hom}, \xi)$.

For $\mathcal{T} = \mathcal{U}_2$, an algebra for the ultrafilter monad $\mathbb{U}$ on $\mathbf{Ord}$ is an *ordered compact Hausdorff space* as introduced in [Nachbin, 1950] (except that we do not assume anti-symmetry here). We recall that these ordered compact Hausdorff spaces are traditionally defined as triples $(X, \leq, \mathcal{O})$ where $(X, \leq)$ is an ordered set and $\mathcal{O}$ is a compact Hausdorff topology on X so that $\{(x, y) \mid x \leq y\}$ is closed in $X \times X$, but the latter requirement means precisely that the convergence $\alpha : UX \rightarrow X$ of $\mathcal{O}$ is monotone (see [Tholen, 2009]). A trivial but important example of an ordered compact Hausdorff space is the two-element chain $2 = \{0, 1\}$ with the discrete topology. We also note that, for an order relation $\leq$ on X ,

$$\mathfrak{x}(U_\xi \leq) \mathfrak{x}' \iff \forall A \in \mathfrak{x}' . \downarrow A \in \mathfrak{x},$$

for all $\mathfrak{x}, \mathfrak{x}' \in UX$.

For $\mathcal{T} = \mathcal{U}_{\mathbf{P}_+}$ it seems natural to call an algebra for the ultrafilter monad $\mathbb{U}$ on $\mathbf{Met}$ a *metric compact Hausdorff space*. The set $[0, \infty]$ equipped with the metric $\text{hom}(u, v) = v \ominus u$ becomes a metric compact Hausdorff space $\mathbf{P}_+$ with the Euclidean compact Hausdorff topology whose convergence is given by $\xi : U[0, \infty] \rightarrow [0, \infty]$ (see Example 1.4 (5)). Similarly, for $\mathcal{T} = \mathcal{U}_{\mathbf{P}_\wedge}$, we call an algebra for the ultrafilter monad $\mathbb{U}$ on $\mathbf{UMet}$ an *ultrametric compact Hausdorff space*, and $[0, \infty]$ becomes an ultrametric compact Hausdorff space where the metric is given by the internal hom of $\mathbf{P}_\wedge$ (see Example 1.4 (4)) and the topology is again the Euclidean compact Hausdorff topology.

There is a canonical functor

$$K : (\mathbf{V}\text{-Cat})^{\mathbb{T}} \rightarrow \mathcal{T}\text{-Cat}.$$

which associates to each $X = (X, a_0, \alpha)$ in $(\mathbf{V}\text{-Cat})^{\mathbb{T}}$ the $\mathcal{T}$ -category $KX = (X, a)$ where $a = a_0 \cdot \alpha$. Note that $(a_0 \cdot \alpha)_0 = a_0$, hence our notation remains consistent. The category $(\mathbf{V}\text{-Cat})^{\mathbb{T}}$ is actually a 2-category with the order relation on hom-sets inherited from $\mathbf{V}\text{-Cat}$, and one easily verifies that K is a 2-functor. Applying K to $\mathbf{V} = (\mathbf{V}, \text{hom}, \xi)$ produces the $\mathcal{T}$ -category $\mathbf{V} = (\mathbf{V}, \text{hom}_\xi)$ where

$$\text{hom}_\xi : TV \times \mathbf{V} \rightarrow \mathbf{V}, (\mathfrak{v}, v) \mapsto \text{hom}(\xi(\mathfrak{v}), v).$$

We note that $\mathbf{V} = (\mathbf{V}, \text{hom}_\xi)$ is separated since $\mathcal{T}\text{-Cat}(G, \mathbf{V}) \simeq \mathbf{V}$ in $\mathbf{Ord}$.

The functor $M : \mathcal{T}\text{-Cat} \rightarrow \mathbf{V}\text{-Cat}$ mentioned at the beginning of this section lifts to a functor $M : \mathcal{T}\text{-Cat} \rightarrow (\mathbf{V}\text{-Cat})^{\mathbb{T}}$ sending (X, a) to $(TX, T_\xi a \cdot m_X^{\circ}, m_X)$. To see this, we have to show that

$$m_X : (TTX, T_\xi(T_\xi a \cdot m_X^{\circ})) \rightarrow (TX, T_\xi a \cdot m_X^{\circ})$$

is a $\mathbf{V}$ -functor. In fact, from $m_X \cdot m_{TX} = m_X \cdot Tm_X$ one obtains $m_{TX} \cdot Tm_X^{\circ} \leq m_X^{\circ} \cdot m_X$ and then

$$m_X \cdot T_\xi T_\xi a \cdot Tm_X^{\circ} \leq T_\xi a \cdot m_{TX} \cdot Tm_X^{\circ} \leq T_\xi a \cdot m_X^{\circ} \cdot m_X.$$

Furthermore, one easily verifies that M is a 2-functor. We denote from now on the “original functor M ” going from $\mathcal{T}\text{-Cat}$ to $\mathbf{V}\text{-Cat}$ as $M_0 : \mathcal{T}\text{-Cat} \rightarrow \mathbf{V}\text{-Cat}$.

Examples 2.2. For a topological space $X = (X, a)$, the order relation $\hat{a} = U_\xi a \cdot m_X^{\circ}$ is given by

$$\mathfrak{x} \hat{a} \mathfrak{y} \quad \text{whenever } \overline{A} \in \mathfrak{y} \text{ for every } A \in \mathfrak{x}.$$

For an approach space $X = (X, a)$, the metric $\hat{a} = U_\xi a \cdot m_X^\circ$ is given by

$$\hat{a}(\mathfrak{x}, \mathfrak{y}) = \inf\{\varepsilon \mid \forall A \in \mathfrak{x} \cdot \overline{A}^{(\varepsilon)} \in \mathfrak{y}\}.$$

Theorem 2.3. $M : \mathcal{T}\text{-Cat} \rightarrow (\mathbf{V}\text{-Cat})^\mathbb{T}$ is left adjoint to $K : (\mathbf{V}\text{-Cat})^\mathbb{T} \rightarrow \mathcal{T}\text{-Cat}$.

Proof. For every $\mathcal{T}$ -category $X = (X, a)$, $e_X : X \rightarrow KM(X)$ is a $\mathcal{T}$ -functor since

$$T_\xi a \cdot m_X^\circ \cdot m_X \cdot Te_X = T_\xi a \cdot m_X^\circ \geq T_\xi a \cdot e_{TX} \geq e_X \cdot a,$$

and we obtain a natural transformation $e : 1 \rightarrow KM$. Let now $X = (X, a_0, \alpha)$ be in $\mathbf{V}\text{-Cat}^\mathbb{T}$. Then α is a $\mathbb{T}$ -algebra homomorphism $\alpha : (TX, m_X) \rightarrow (X, \alpha)$, and also a $\mathbf{V}$ -functor $\alpha : (TX, T_\xi(a_0 \cdot \alpha) \cdot m_X^\circ) \rightarrow (X, a_0)$ since

$$\alpha \cdot T_\xi(a_0 \cdot \alpha) \cdot m_X^\circ = \alpha \cdot T_\xi(a_0) \cdot T\alpha \cdot m_X^\circ \leq a_0 \cdot \alpha \cdot T\alpha \cdot m_X^\circ = a_0 \cdot \alpha.$$

Clearly, for every $f : X \rightarrow Y$ in $\mathbf{V}\text{-Cat}^\mathbb{T}$ where $X = (X, a_0, \alpha)$ and $Y = (Y, a_0, \beta)$, the diagram

$$\begin{array}{ccc} MK(X) & \xrightarrow{\alpha} & X \\ Tf=MK(f) \downarrow & & \downarrow f \\ MK(Y) & \xrightarrow{\beta} & Y \end{array}$$

commutes, hence the family $(\alpha)_{(X, a_0, \alpha)}$ is a natural transformation $MK \rightarrow 1$. Finally, for (X, a) in $\mathcal{T}\text{-Cat}$ and $(Y, b_0, \beta) \in \mathbf{V}\text{-Cat}^\mathbb{T}$,

$$m_X \cdot Me_X = 1_{MX} \quad \text{and} \quad \beta \cdot e_Y = 1_Y;$$

and the assertion follows. $\square$

Via the adjunction $M \dashv K$ one obtains a lifting of the **Set**-monad $\mathbb{T} = (T, e, m)$ to a monad on $\mathcal{T}\text{-Cat}$, also denoted as $\mathbb{T} = (T, e, m)$. Explicitly, $T : \mathcal{T}\text{-Cat} \rightarrow \mathcal{T}\text{-Cat}$ sends a $\mathcal{T}$ -category $X = (X, a)$ to $(TX, T_\xi a \cdot m_X^\circ \cdot m_X)$. Moreover, $e_X : X \rightarrow TX$ is fully faithful since $e_X^\circ \cdot T_\xi a \cdot m_X^\circ \cdot m_X \cdot Te_X = a$.

Proposition 2.4. *The monad $\mathbb{T} = (T, e, m)$ on $\mathcal{T}\text{-Cat}$ is of Kock-Zöberlein type.*

Proof. We show that $m_X \dashv e_{TX}$ in $\mathcal{T}\text{-Cat}$. By definition, $m_X \cdot e_{TX} = 1_{TX}$. To see $k \leq s(\mathfrak{X}, e_{TX} \cdot m_X(\mathfrak{X}))$ for all $\mathfrak{X} \in TTX$, we show $m_X^\circ \cdot e_{TX}^\circ \cdot s \geq 1_{TTX}$ (where $s = T_\xi(r \cdot m_X) \cdot m_{TX}^\circ$ and $r = T_\xi a \cdot m_X^\circ$) in $\mathbf{V}\text{-Rel}$. In fact,

$$m_X^\circ \cdot e_{TX}^\circ \cdot T_\xi(r \cdot m_X) \cdot m_{TX}^\circ = m_X^\circ \cdot r \cdot m_X \geq m_X^\circ \cdot m_X \geq 1_{TTX}. \quad \square$$

Since the monad $\mathbb{T}$ on $\mathcal{T}\text{-Cat}$ is of Kock-Zöberlein type, an algebra structure $\alpha : TX \rightarrow X$ on a $\mathcal{T}$ -category X is left adjoint to the unit $e_X : X \rightarrow TX$. However, unless X is separated, a left adjoint $\alpha : TX \rightarrow X$ to e_X is in general only a pseudo-algebra structure on X , that is,

$$\alpha \cdot e_X \simeq 1_X \quad \text{and} \quad \alpha \cdot T\alpha \simeq \alpha \cdot m_X. \quad (2)$$

Definition 2.5. We call a $\mathcal{T}$ -category X *representable* whenever $e_X : X \rightarrow TX$ has a left adjoint in $\mathcal{T}\text{-Cat}$. A $\mathcal{T}$ -functor $f : X \rightarrow Y$ between representable $\mathcal{T}$ -categories X and Y , with left adjoint $\alpha : TX \rightarrow X$ and $\beta : TY \rightarrow Y$ respectively, is called a *pseudo-homomorphism* whenever

$$\beta \cdot Tf \simeq f \cdot \alpha.$$

Of course, if Y is separated, then one has equality above. We denote the category of representable $\mathcal{T}$ -categories and pseudo-homomorphism by

$$\mathcal{T}\text{-ReprCat},$$

and its full subcategory defined by the separated representable $\mathcal{T}$ -categories by $\mathcal{T}\text{-ReprCat}_{\text{sep}}$.

Remark 2.6. We borrowed the designation “representable” from [Hermida, 2000] where the notion of representable multicategory via a “monadic 2-adjunction between the 2-category of strict monoidal categories and that of multicategories” is introduced. In a nutshell, strict monoidal categories are to multicategories what ordered compact Hausdorff spaces are to topological spaces.

Remark 2.7. For a separated representable $\mathcal{T}$ -category $X = (X, a)$, the left adjoint $\alpha : TX \rightarrow X$ to $e_X : X \rightarrow TX$ is unique and actually the structure of a $\mathbb{T}$ -algebra on X . Therefore there is a canonical forgetful functor $\mathcal{T}\text{-ReprCat}_{\text{sep}} \rightarrow \text{Set}^{\mathbb{T}}$ sending (X, a) to (X, α) which is part of an adjunction

$$\mathcal{T}\text{-ReprCat}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad} \\ \top \\ \xleftarrow{\quad} \end{array} \text{Set}^{\mathbb{T}}$$

where the left adjoint $\text{Set}^{\mathbb{T}} \rightarrow \mathcal{T}\text{-ReprCat}_{\text{sep}}$ interprets the $\mathbb{T}$ -structure α on a set X as a $\mathcal{T}$ -structure on X .

Proposition 2.8. *The following assertions are equivalent, for a $\mathcal{T}$ -category $X = (X, a)$.*

- (i) *X is representable.*
- (ii) *X is core-compact and there is a map $\alpha : TX \rightarrow X$ such that $a = a_0 \cdot \alpha$.*

Proof. Assume first that X is representable. Then $e_X : X \rightarrow TX$ has a left adjoint $\alpha : TX \rightarrow X$ in $\mathcal{T}\text{-Cat}$ which necessarily satisfies (2). Hence also $\alpha \dashv e_X$ in $\mathbf{V}\text{-Cat}$, which gives (with $\hat{a} = T_{\xi} \cdot m_X^{\circ}$)

$$a(\mathfrak{x}, x) = \hat{a}(\mathfrak{x}, e_X(x)) = a_0(\alpha(\mathfrak{x}), x),$$

and then we calculate $a_0 \cdot \alpha \cdot m_X = a_0 \cdot \alpha \cdot T\alpha \leq a_0 \cdot \alpha \cdot T_{\xi}(a_0 \cdot \alpha)$. Conversely, assume now (ii). Then $e_X^{\circledast} = e_X^{\circ} \cdot T_{\xi}a \cdot m_X^{\circ} \cdot m_X = a \cdot m_X$ and $\alpha_{\circledast} = a \cdot T\alpha = a \cdot T_{\xi}a_0 \cdot T\alpha = a \cdot T_{\xi}a = a \cdot m_X$, hence α is a $\mathcal{T}$ -functor and $\alpha \dashv e_X$ in $\mathcal{T}\text{-Cat}$. $\square$

The following result is easy to prove.

Lemma 2.9. *Let (X, a) , (Y, b) be representable $\mathcal{T}$ -categories with left adjoints $\alpha : TX \rightarrow X$ and $\beta : TY \rightarrow Y$ respectively, and let $f : X \rightarrow Y$ be a map. Then f is a $\mathcal{T}$ -functor $f : (X, a) \rightarrow (Y, b)$ if and only if $f : (X, a_0) \rightarrow (Y, b_0)$ is a $\mathbf{V}$ -functor and $\beta \cdot Tf(\mathfrak{x}) \leq f \cdot \alpha(\mathfrak{x})$, for all $\mathfrak{x} \in TX$.*

We have the comparison functor

$$K^{\mathbb{T}} : (\mathbf{V}\text{-Cat})^{\mathbb{T}} \rightarrow (\mathcal{T}\text{-Cat})^{\mathbb{T}}$$

which sends (X, a_0, α) in $(\mathbf{V}\text{-Cat})^{\mathbb{T}}$ to $(X, a_0 \cdot \alpha, \alpha)$. Furthermore, the forgetful functor $(-)_0 : \mathcal{T}\text{-Cat} \rightarrow \mathbf{V}\text{-Cat}$ lifts to a functor $(-)_0^{\mathbb{T}} : (\mathcal{T}\text{-Cat})^{\mathbb{T}} \rightarrow (\mathbf{V}\text{-Cat})^{\mathbb{T}}$ since the identity map $T(X_0) \rightarrow (TX)_0$ is a $\mathbf{V}$ -functor, for each $\mathcal{T}$ -category X . Clearly, $(K^{\mathbb{T}}X)_0^{\mathbb{T}} = X$ for every X in $(\mathbf{V}\text{-Cat})^{\mathbb{T}}$, and for $X = (X, a)$ in $(\mathcal{T}\text{-Cat})^{\mathbb{T}}$ with Eilenberg–Moore structure $\alpha : TX \rightarrow X$ one has $a = a_0 \cdot \alpha$ by Proposition 2.8. We conclude:

Theorem 2.10. $(\mathbf{V}\text{-Cat})^{\mathbb{T}} \simeq (\mathcal{T}\text{-Cat})^{\mathbb{T}}$.

The notion of pseudo-algebra was already lurking in the discussion above. In fact, just as for algebras, every pseudo-algebra structure $\alpha : TX \rightarrow X$ on a $\mathbf{V}$ -category $X = (X, a_0)$ gives rise to the representable $\mathcal{T}$ -category $(X, a_0 \cdot \alpha)$, and equivalent pseudo-algebra structures induce the same $\mathcal{T}$ -category. Moreover, by Proposition 2.8, every representable $\mathcal{T}$ -category is of this form.

Our next aim is to introduce a concept of dual $\mathcal{T}$ -category which generalises the one for $\mathbf{V}$ -categories.

Definition 2.11. A $\mathcal{T}$ -graph $X = (X, a)$ is called *dualisable* whenever $a_0 = a \cdot e_X$ is transitive and $a = a_0 \cdot \alpha$, for some map $\alpha : TX \rightarrow X$.

Every $\mathcal{T}$ -category (X, a) where $a = a_0 \cdot \alpha$ (for some map $\alpha : TX \rightarrow X$) is a dualisable $\mathcal{T}$ -graph. Another important example will be provided by Lemma 4.1. For a dualisable $\mathcal{T}$ -graph $X = (X, a)$, we write X_0 to denote its underlying $\mathbf{V}$ -category $X_0 = (X, a_0)$. We consider TX as a discrete $\mathbf{V}$ -category, so that $\alpha : TX \rightarrow X_0$ is a $\mathbf{V}$ -functor. With this notation, $a_0 \cdot \alpha = \alpha_*$ and, if $\alpha_* = a = \beta_*$, also $\alpha^* = \beta^*$ and therefore

$$a_0^\circ \cdot \alpha = (\alpha^*)^\circ = (\beta^*)^\circ = a_0^\circ \cdot \beta.$$

Lemma 2.12. *Let $X = (X, a)$ be a dualisable $\mathcal{T}$ -graph. Then $(X, a_0^\circ \cdot \alpha)$ is a dualisable $\mathcal{T}$ -graph as well, and the underlying $\mathbf{V}$ -category of $(X, a_0^\circ \cdot \alpha)$ is $(X_0)^{\text{op}}$.*

Proof. It suffices to show $a_0^\circ = a_0^\circ \cdot \alpha \cdot e_X$. From $a = a_0 \cdot \alpha$ we infer $a_0 = a_0 \cdot \alpha \cdot e_X = (\alpha \cdot e_X)_*$, hence $a_0 = (\alpha \cdot e_X)^*$ and therefore $a_0^\circ = a_0^\circ \cdot \alpha \cdot e_X$. $\square$

Definition 2.13. Let $X = (X, a)$ be a dualisable $\mathcal{T}$ -graph. Then the *dual* $\mathcal{T}$ -graph X^{op} of X is $X^{\text{op}} = (X, a_0^\circ \cdot \alpha)$.

By the discussion above, this definition is independent of the choice of α . Of course, the dual of a $\mathbf{V}$ -category in the sense above is just the usual dual. Also note that, even if X is a $\mathcal{T}$ -category, X^{op} need not be a $\mathcal{T}$ -category (see Proposition 2.15 below).

The following result is a variation of Lemma 2.9.

Lemma 2.14. *For a $(\mathbb{T}, \mathbf{V})$ -functor $f : (X, a) \rightarrow (Y, b)$ between dualisable $(\mathbb{T}, \mathbf{V})$ -graphs with $a = a_0 \cdot \alpha$ and $b = b_0 \cdot \beta$, the map $f : X \rightarrow Y$ also defines a $(\mathbb{T}, \mathbf{V})$ -functor $f^{\text{op}} : X^{\text{op}} \rightarrow Y^{\text{op}}$ if and only if $f \cdot \alpha \simeq \beta \cdot Tf$.*

Proposition 2.15. *Let $X = (X, a)$ be a $\mathcal{T}$ -category where $a = a_0 \cdot \alpha$, for some map $\alpha : TX \rightarrow X$. Then the following assertions are equivalent.*

- (i) *The $\mathcal{T}$ -graph X^{op} is actually a $\mathcal{T}$ -category.*
- (ii) *X is core-compact.*
- (iii) *X is representable.*

Proof. Proposition 2.8 affirms (ii) $\Leftrightarrow$ (iii), and (iii) $\Rightarrow$ (i) is clear. Assume now that X^{op} is a $\mathcal{T}$ -category. Since X is a $\mathcal{T}$ -category,

$$(\alpha \cdot T\alpha)_* = a_0 \cdot \alpha \cdot T\alpha \leq a_0 \cdot \alpha \cdot T_\xi a_0 \cdot T\alpha \leq a_0 \cdot \alpha \cdot m_X = (a_0 \cdot \alpha)_*;$$

similarly, since X^{op} is a $\mathcal{T}$ -category,

$$a_0^\circ \cdot \alpha \cdot T\alpha \leq a_0^\circ \cdot \alpha \cdot m_X$$

and therefore $(\alpha \cdot T\alpha)^* \leq (a_0 \cdot \alpha)^*$. Consequently, $(\alpha \cdot T\alpha)_* = (a_0 \cdot \alpha)_*$, hence $a \cdot T_\xi a = a \cdot m_X$. $\square$

In conclusion, taking duals gives a functor $(-)^{\text{op}} : \mathcal{T}\text{-ReprCat} \rightarrow \mathcal{T}\text{-ReprCat}$ which makes the diagram

$$\begin{array}{ccc} \mathcal{T}\text{-ReprCat} & \xrightarrow{(-)^{\text{op}}} & \mathcal{T}\text{-ReprCat} \\ (-)_0 \downarrow & & \downarrow (-)_0 \\ \mathbf{V}\text{-Cat} & \xrightarrow{(-)^{\text{op}}} & \mathbf{V}\text{-Cat} \end{array}$$

commutative.

By Propositions 1.7 and 2.8, every representable $\mathcal{T}$ -category is $\otimes$ -exponentiable. It is interesting to observe that the canonical map $Y^{(X, a_0 \cdot \alpha)} \rightarrow Y^{(X, \alpha)}$ is actually an embedding, for every (X, a_0, α) in $(\mathbf{V}\text{-Cat})^{\mathbb{T}}$ (see [Hofmann, 2010, Lemma 5.2] for a proof for the approach case, the general case is similar). For a $\mathcal{T}$ -category X , its *presheaf $\mathcal{T}$ -category* PX is defined as $PX := \mathbf{V}^{(TX)^{\text{op}}}$ with structure relation denoted as $\llbracket -, - \rrbracket$. By the observation above, this definition coincides with the one given in [Clementino and Hofmann, 2009a]. Proposition 1.7

implies that the underlying $\mathbf{V}$ -category $(PX)_0$ is a full subcategory of the presheaf $\mathbf{V}$ -category of $M_0(X)$, where, for $\psi, \psi' \in PX$,

$$[\psi, \psi'] := \llbracket e_{PX}(\psi), \psi' \rrbracket = \bigwedge_{\mathfrak{x} \in TX} \text{hom}(\psi(\mathfrak{x}), \psi'(\mathfrak{x})).$$

A slight adaptation of [Clementino and Hofmann, 2009a, Theorem 2.5] gives

Theorem 2.16. *Let $\varphi : X \multimap Y$ be a $\mathcal{T}$ -relation. The following assertions are equivalent.*

- (i) $\varphi : X \multimap Y$ is a $\mathcal{T}$ -distributor.
- (ii) $\varphi : (TX)^{\text{op}} \otimes Y \rightarrow \mathbf{V}$ is a $\mathcal{T}$ -functor.
- (iii) $\ulcorner \varphi \urcorner : Y \rightarrow PX$ is a $\mathcal{T}$ -functor.

For each $\mathcal{T}$ -category $X = (X, a)$, $a : X \multimap X$ is a $\mathcal{T}$ -distributor which gives us the *Yoneda functor*

$$y_X = \ulcorner a \urcorner : X \rightarrow PX.$$

We recall the following result from [Hofmann, 2011].

Theorem 2.17. *Let $\psi : X \multimap Z$ and $\varphi : X \multimap Y$ be $\mathcal{T}$ -distributors. Then, for all $\mathfrak{z} \in TZ$ and $y \in Y$,*

$$\llbracket T \ulcorner \psi \urcorner(\mathfrak{z}), \ulcorner \varphi \urcorner(y) \rrbracket = (\varphi \circ \psi)(\mathfrak{z}, y).$$

In particular, for each $\psi \in PX$ and each $\mathfrak{x} \in TX$, $\psi(\mathfrak{x}) = \llbracket T y_X(\mathfrak{x}), \psi \rrbracket$.

Example: topological (and approach) spaces. Regarding $\mathcal{T} = \mathcal{U}_2$, an ordered compact Hausdorff space $X = (X, \leq, \alpha)$ induces a topological space X by stipulating that an ultrafilter $\mathfrak{x} \in UX$ converges to $x \in X$ whenever $\alpha(\mathfrak{x}) \leq x$, that is, by making $\alpha(\mathfrak{x})$ the smallest convergence point of $\mathfrak{x}$. The ordered compact Hausdorff space 2 (see Example 2.1) induces the Sierpiński space 2 where $\{1\}$ is closed and $\{0\}$ is open, and 2^{op} has $\{1\}$ open and $\{0\}$ closed. Representable topological T_0 -spaces (under the name *stably compact spaces* or *well-compact spaces*) are well studied, we refer to [Simmons, 1982; Jung, 2004; Lawson, 2011] for more information. Below and until the end of this section we develop some well-known basic properties of these spaces, mainly to connect the convergence-theoretic perspective of this paper with the classical account via open subsets.

By Proposition 2.8, every representable topological space $X = (X, a)$ satisfies $a \cdot U_\xi a = a \cdot m_X$, which is equivalent to X being core-compact (see [Pisani, 1999]). In fact, slightly more can be said:

Lemma 2.18. *Every representable topological space is locally compact.*

Proof. For every topological space X , the topology on UX is generated by all sets of the form

$$A^\# = \{\mathfrak{a} \in UX \mid A \in \mathfrak{a}\},$$

where $A \subseteq X$ is open. Furthermore, for any ultrafilter $\mathfrak{X} \in UUX$ with $A^\# \in \mathfrak{X}$, one has $m_X(\mathfrak{X}) \in A^\#$ and therefore $A^\#$ is compact; hence UX is locally compact. If X is representable, then X is a split subobject of UX (since $\alpha : UX \rightarrow X$ can be chosen so that $\alpha(e_X(x)) = x$) and therefore also locally compact. $\square$

Hence, a topological space X is representable if and only if X is locally compact and every ultrafilter on X has a smallest convergence point (see Proposition 2.8). The latter condition says in particular that the set of limit points of an ultrafilter $\mathfrak{x}$ is irreducible. Since in any topological space an irreducible closed subset is the set of limit points of some ultrafilter, we find that X is representable if and only if X is locally compact, weakly sober (every irreducible closed subset is the closure of some point) and, for every $\mathfrak{x} \in UX$, the set of limit points of $\mathfrak{x}$ is irreducible. For any core-compact topological space X , the last condition is equivalent to stability of the way-below relation on the lattice of open subsets under finite intersections: $\bigcap_i U_i \ll \bigcap_i V_i$, for open subsets $U_1, \dots, U_n$ and $V_1, \dots, V_n$ ($n \in \mathbb{N}$) of X with $U_i \ll V_i$ for each $1 \leq i \leq n$ (see

[Simmons, 1982]). By definition, $U \ll V$ whenever every open cover of V contains a finite subcover of U , which is the case if and only if every ultrafilter $\mathfrak{x}$ with $U \in \mathfrak{x}$ has a limit point in V . If X is representable, $U \ll V$ if and only if any smallest limit point of an ultrafilter $\mathfrak{x}$ with $U \in \mathfrak{x}$ belongs to V . Hence:

Proposition 2.19. *A topological space X is representable if and only if X is locally compact, weakly sober and the way-below relation on the lattice of opens is stable under finite intersection.*

This stability condition on the way-below relation is sometimes replaced by a stability condition on the compact down-sets of X , as we explain next. Clearly, for X representable, $\bigcap \emptyset = X$ is compact, and binary intersections of pairs of compact down-sets are compact: if $A, B \subseteq X$ are compact down-sets and $A \cap B \in \mathfrak{x}$, then any smallest convergence point of $\mathfrak{x}$ belongs to both A and B and therefore also to $A \cap B$. Secondly, since open subsets are down-closed, the down-closure (with respect to the underlying order) of a compact subset of a topological space is compact. Therefore, for a locally compact space X and $U, V \subseteq X$ open,

$$U \ll V \iff U \subseteq K \subseteq V \text{ for some compact down-set } K \subseteq X. \quad (3)$$

From that one sees at once that stability of the way-below relation under finite intersection follows from stability of compact down-sets under finite intersection.

Proposition 2.20. *A topological space X is representable if and only if X is locally compact, weakly sober and finite intersections of compact down-sets are compact.*

By definition, a pseudo-homomorphism between representable topological spaces is a continuous map $f : X \rightarrow Y$ which preserves smallest convergence points of ultrafilters.

Proposition 2.21. *Let $f : X \rightarrow Y$ be a continuous map between representable topological spaces. Then the following assertions are equivalent.*

- (i) f is a pseudo-homomorphism.
- (ii) For every compact down-set $K \subseteq Y$, $f^{-1}(K)$ is compact.
- (iii) The frame homomorphism $f^{-1} : \mathcal{O}Y \rightarrow \mathcal{O}X$ preserves the way-below relation.

Proof. Clearly, (i) $\Rightarrow$ (ii); and the implication (ii) $\Rightarrow$ (iii) follows from (3). Assume now (iii) and let $x \in X$ be a smallest convergence point of $\mathfrak{x} \in UX$. Assume that $Uf(\mathfrak{x}) \rightarrow y \in Y$. Let $U, V \subseteq Y$ be open subsets with $y \in U \ll V$. Then $f^{-1}(U) \ll f^{-1}(V)$ and $f^{-1}(U) \in \mathfrak{x}$, hence $x \in f^{-1}(V)$ and therefore $f(x) \in V$. We conclude that $f(x) \leq y$. $\square$

A continuous map $f : X \rightarrow Y$ satisfying condition (ii) (and hence also (i) and (iii)) above is called *spectral*.

Corollary 2.22. *Let X be a representable topological space. A continuous map $\varphi : X \rightarrow 2$ is a homomorphism if and only if the open set $\varphi^{-1}(0)$ is compact.*

From Lemma 2.9 we obtain

Proposition 2.23. *Let $(X, \leq, \alpha)$ be an ordered compact Hausdorff space and let $a = \leq \cdot \alpha$ be the induced topology. A subset $A \subseteq X$ is open in (X, a) if and only if A is down-closed and open in the compact Hausdorff space (X, α) .*

Proposition 2.24. *Let X be a representable space, $\mathfrak{x} \in UX$ and $x_0 \in X$ be a smallest convergence point of $\mathfrak{x}$. For any $x \in X$, $x \leq x_0$ if and only if $\mathfrak{x}$ contains all complements of compact down-sets B with $x \notin B$.*

Proof. If $x \leq x_0$, then $\mathfrak{x}$ cannot contain any compact down-sets B with $x \notin B$. Assume now that $\mathfrak{x}$ contains these subsets. Take a neighbourhood B of x_0 where B is a compact down-set. Then $x \in B$ since otherwise $B \in \mathfrak{x}$ and $X \setminus B \in \mathfrak{x}$. $\square$

Corollary 2.25. *Let X be a representable space. Then the topology of X^{op} is generated by the complements of compact down-sets B of X . Furthermore, the ultrafilter convergence of the topology generated by the opens and the complements of compact down-sets of X is given by taking smallest convergence points of ultrafilters of X .*

We note that this notion of dual space was introduced by M. Hochster (see [Hochster, 1969]).

Corollary 2.26. *Let $(X, \leq, \alpha)$ be an anti-symmetric ordered compact Hausdorff space. Then the topology of (X, α) is generated by the open subsets and the complements of compact down-sets of the representable space $(X, \leq \cdot \alpha)$.*

It is “folklore” that the category of anti-symmetric ordered compact Hausdorff spaces and homomorphisms is equivalent to the category **StablyComp** of stably compact spaces and spectral maps (the first appearance of this result seems to be [Gierz *et al.*, 1980]), which is the restriction of Theorem 2.10 to separated spaces. A stably compact space X is called *spectral* whenever the compact opens form a basis for the topology (which is equivalent to the statement that the source $(\varphi : X \rightarrow 2)$ of all homomorphisms into 2 is point-separating and initial in **Top**, and also in the category of stably compact spaces and spectral maps). For a spectral space X and $U, V \subseteq X$ open, $U \ll V$ if and only if $U \subseteq K \subseteq V$ for some compact open subset $K \subseteq X$; hence a continuous map $f : Y \rightarrow X$ (where Y is representable) is spectral if and only if the inverse image of every compact open subset of X is compact in Y . A famous result of M.H. Stone [Stone, 1938] states that the category of spectral spaces and spectral maps is dually equivalent to the category **DLat** of distributive lattices and homomorphisms. A different perspective on this duality was given in [Priestley, 1970]: **DLat** is also dually equivalent to the category of (nowadays called) *Priestley spaces* and homomorphisms. Here a Priestley space is an anti-symmetric ordered compact Hausdorff space where $x \not\leq y$ implies the existence of an clopen down-set V and a clopen up-set U with $x \in U$, $y \in V$ and $U \cap V = \emptyset$ (equivalently: the source $(\varphi : X \rightarrow 2)$ of all homomorphisms into 2 is point-separating and initial in the category of ordered compact Hausdorff spaces and homomorphisms). In particular, both results together imply the equivalence between spectral spaces and Priestley spaces which is a restriction of the aforementioned equivalence between stably compact spaces and anti-symmetric ordered compact Hausdorff spaces. We also note that, for X compact Hausdorff, X is spectral if and only if the simultaneously closed and open subsets of X form a basis for the topology of X , i.e. if X is a *Stone space*. Every continuous map between Stone spaces is spectral, and the full subcategory **Stone** of **Spec** defined by all Stone spaces is dually equivalent to the category **Bool** of Boolean algebras and homomorphisms ([Stone, 1936; Johnstone, 1986]).

The case of metric compact Hausdorff spaces (we consider now $\mathcal{T} = \mathcal{U}_{\text{P}_+}$) was studied in [Gutierrez and Hofmann, 2012]). An approach space $X = (X, a)$ is representable if and only if X is weakly sober, $+$ -exponentiable and has the property that $a(\mathfrak{x}, -)$ is an approach prime element, for every $\mathfrak{x} \in UX$ (we refer to [Banaschewski *et al.*, 2006] and [Van Olmen, 2005] for the theory of sober approach space). The metric compact Hausdorff space P_+ (see Example 2.1) induces the “Sierpiński approach space” P_+ with approach convergence structure $\lambda(\mathfrak{x}, x) = x \ominus \xi(\mathfrak{x})$; but, in contrast to the topological case, P_+ is not isomorphic to P_+^{op} (for instance, P_+ is injective but P_+^{op} is not). Similarly, the ultrametric compact Hausdorff space $\text{P}_\wedge$ produces the approach space $\text{P}_\wedge$. We note that both P_+ and $\text{P}_\wedge$ have the same underlying topological space.

3. COCOMPLETE $\mathcal{T}$ -CATEGORIES

By an appropriate translation from the **V** to the $\mathcal{T}$ -case one can transport the notions of weighted colimit (see [Eilenberg and Kelly, 1966; Kelly, 1982]) and cocompleteness into the realm $\mathcal{T}$ -categories, as we recall now briefly from [Hofmann, 2011] and [Clementino and Hofmann, 2009b]. A weighted colimit diagram in a $\mathcal{T}$ -category X is given by a $\mathcal{T}$ -functor $d : D \rightarrow X$ and

a $\mathcal{T}$ -distributor $\varphi : D \multimap G$ (where $G = (1, e_1^\circ)$).

$$\begin{array}{ccc} D & \xrightarrow{d} & X \\ \varphi \circ & \downarrow & \\ & G & \end{array}$$

A *colimit* of such weighted diagram is an element $x \in X$ which represents $d_\circ \multimap \varphi$, that is, $x_\circ = d_\circ \multimap \varphi$. If such x exists, it is unique up to equivalence, and one calls x a φ -*weighted colimit* of d and writes $x \simeq \text{colim}(d, \varphi)$. We say that a $\mathcal{T}$ -functor $f : X \rightarrow Y$ *preserves* the φ -weighted colimit x of d if $f(x)$ is the φ -weighted colimit of $f \cdot d$, that is, if $f(x)_\circ = (f \cdot d)_\circ \multimap \varphi$. A $\mathcal{T}$ -functor $f : X \rightarrow Y$ is called *cocontinuous* if it preserves all weighted colimits which exist in X , and a $\mathcal{T}$ -category X is *cocomplete* if every weighted colimit diagram in X has a colimit in X . As in the $\mathbf{V}$ -category case, cocompleteness of X follows from the existence of colimits along identities. In fact, for any weight $\varphi : D \multimap G$, the φ -weighted colimit of d exists if and only if the $(\varphi \circ d^\circ)$ -weighted colimit of $1_X : X \rightarrow X$ exists, and in that case one has $\text{colim}(d, \varphi) \simeq \text{colim}(1_X, \varphi \circ d^\circ)$. Moreover, a $\mathcal{T}$ -functor $f : X \rightarrow Y$ preserves the φ -weighted colimit of d if and only if it preserves the $(\varphi \circ d^\circ)$ -weighted colimit of 1_X . In the sequel we will write $\text{Sup}_X(\psi)$ (or simply $\text{Sup}(\psi)$) instead of $\text{colim}(1_X, \psi)$.

For a cocomplete $\mathcal{T}$ -category X , the map $\text{Sup}_X : PX \rightarrow X$ turns out to be left adjoint to the Yoneda embedding $y_X : X \rightarrow PX$ in $\mathbf{V}\text{-Cat}$; however, Sup_X is in general not a $\mathcal{T}$ -functor (see [Hofmann and Waszkiewicz, 2011, Example 5.7]). A $\mathcal{T}$ -category X is called *totally cocomplete* whenever $y_X : X \rightarrow PX$ has a left adjoint $\text{Sup}_X : PX \rightarrow X$ in $\mathcal{T}\text{-Cat}$. Curiously, total cocompleteness can be characterised by the existence of a slightly more general type of colimits, as we explain next. From now on we let in a *weighted colimit diagram* the weight $\varphi : D \multimap A$ be an arbitrary $\mathcal{T}$ -distributor. A colimit of such a diagram is a $\mathcal{T}$ -functor $g : A \rightarrow X$ which represents $d_\circ \multimap \varphi$ in the sense that $g_\circ = d_\circ \multimap \varphi$, and we write $g \simeq \text{colim}(d, \varphi)$. We note that one still has $\text{colim}(d, \varphi) \simeq \text{colim}(1_X, \varphi \circ d^\circ)$. A $\mathcal{T}$ -functor $f : X \rightarrow Y$ preserves the φ -weighted colimit g of d if $f \cdot g$ is the φ -weighted colimit of $f \cdot d$, that is, if $(f \cdot g)_\circ = (f \cdot d)_\circ \multimap \varphi$. We write

$\mathcal{T}\text{-CoCts}$

to denote the category of totally cocomplete $\mathcal{T}$ -categories and weighted colimit preserving $\mathcal{T}$ -functors, and $\mathcal{T}\text{-CoCts}_{\text{sep}}$ for its full subcategory defined by the separated $\mathcal{T}$ -categories. As before, in the $\mathbf{V}$ -case we use the designations $\mathbf{V}\text{-CoCts}$ and $\mathbf{V}\text{-CoCts}_{\text{sep}}$.

For every $\mathcal{T}$ -distributor $\varphi : X \multimap Y$, the function $- \circ \varphi : PY \rightarrow PX$ is actually a $\mathcal{T}$ -functor $P\varphi : PY \rightarrow PX$, and this construction yields a functor $P : \mathcal{T}\text{-Dist}^{\text{op}} \rightarrow \mathcal{T}\text{-Cat}$. In fact:

Theorem 3.1. *The functor $P : \mathcal{T}\text{-Dist}^{\text{op}} \rightarrow \mathcal{T}\text{-Cat}$ is right adjoint to $(-)^{\circ} : \mathcal{T}\text{-Cat} \rightarrow \mathcal{T}\text{-Dist}^{\text{op}}$. The units of this adjunction are given by $y_X : X \rightarrow PX$ and $(y_X)_\circ : X \multimap PX$ respectively. The induced monad $\mathbb{P} = (P, y, m)$ on $\mathcal{T}\text{-Cat}$ is of Kock-Zöberlein type (here $m_X = - \circ (y_X)_\circ$).*

Theorem 3.2. *The following assertions are equivalent, for a $\mathcal{T}$ -category X .*

- (i) X is injective w.r.t. fully faithful $\mathcal{T}$ -functors.
- (ii) $y_X : X \rightarrow PX$ has a left inverse $\text{Sup}_X : PX \rightarrow X$, that is, $\text{Sup}_X \cdot y_X \simeq 1_X$.
- (iii) $y_X : X \rightarrow PX$ has a left adjoint $\text{Sup}_X : PX \rightarrow X$.
- (iv) X has all weighted colimits (in the generalised sense).

Here a $\mathcal{T}$ -category X is called *injective* if, for all $\mathcal{T}$ -functors $f : A \rightarrow X$ and fully faithful $\mathcal{T}$ -functors $i : A \rightarrow B$, there exists a $\mathcal{T}$ -functor $g : B \rightarrow X$ such that $g \cdot i \simeq f$. Clearly, for a separated $\mathcal{T}$ -category X we have then $g \cdot i = f$.

Remark 3.3. In the proof of (ii) $\Rightarrow$ (iii) one shows that any left inverse of $y_X : X \rightarrow PX$ is actually left adjoint to y_X . Then, given a left adjoint $\text{Sup}_X : PX \rightarrow X$ of y_X , the colimit of a diagram defined by $d : D \rightarrow X$ and $\varphi : D \multimap A$ can be calculated as $\text{Sup}_X \cdot Pd \cdot \ulcorner \varphi \urcorner$.

Corollary 3.4. *For each $\mathcal{T}$ -category X , PX is cocomplete where $\text{Sup}_{PX} = - \circ (y_X)_\otimes$.*

Proposition 3.5. *Let $f : X \rightarrow Y$ be a $\mathcal{T}$ -functor between totally cocomplete $\mathcal{T}$ -categories. Then the following assertions are equivalent.*

- (i) *f preserves all weighted colimits (in the generalised sense).*
- (ii) *f preserves all weighted colimits with weight of type $D \multimap G$.*
- (iii) *The diagram*

$$\begin{array}{ccc} PX & \xrightarrow{Pf} & PY \\ \text{Sup}_X \downarrow & \simeq & \downarrow \text{Sup}_Y \\ X & \xrightarrow{f} & Y \end{array}$$

commutes up to equivalence.

Theorem 3.6. *The category $\mathcal{T}\text{-CoCts}_{\text{sep}}$ is precisely the category $(\mathcal{T}\text{-Cat})^{\mathbb{P}}$ of Eilenberg–Moore algebras for $\mathbb{P}$. Moreover, the canonical forgetful functors $\mathcal{T}\text{-CoCts}_{\text{sep}} \rightarrow \mathbf{V}\text{-Cat}$ and $\mathcal{T}\text{-CoCts}_{\text{sep}} \rightarrow \mathbf{Set}$ are both monadic.*

Proposition 3.7. *Every left adjoint $\mathcal{T}$ -functor is cocontinuous. A $\mathcal{T}$ -functor between cocomplete $\mathcal{T}$ -categories is left adjoint if and only if it is cocontinuous.*

For a $\mathcal{T}$ -category $X = (X, a)$, one has the $\mathbf{V}$ -category structure $\hat{a} := T_\xi a \cdot m_X^\circ : TX \multimap TX$ on TX , which is indeed a $\mathcal{T}$ -distributor $\hat{a} : X \multimap TX$ since

$$\begin{aligned} \hat{a} \cdot m_X \cdot T_\xi \hat{a} \cdot m_X^\circ &\leq \hat{a} \cdot \hat{a} \cdot m_X \cdot m_X^\circ = \hat{a}, \\ \hat{a} \cdot T_\xi a \cdot m_X^\circ &= \hat{a} \cdot \hat{a} = \hat{a}. \end{aligned}$$

Therefore one obtains a $\mathcal{T}$ -functor $\mathcal{Y}_X : TX \rightarrow PX$, and one easily verifies $\mathcal{Y}_X \cdot e_X = y_X$. Consequently:

Proposition 3.8. *Every totally cocomplete $\mathcal{T}$ -category is representable. Moreover, every left adjoint $\mathcal{T}$ -functor between representable $\mathcal{T}$ -categories is a pseudo-homomorphism.*

Remark 3.9. From the proposition above we obtain a forgetful functor $\mathcal{T}\text{-CoCts} \rightarrow \mathcal{T}\text{-ReprCat}$ which restricts to separated objects. Therefore we also have functors (see Remark 2.7)

$$\mathcal{T}\text{-CoCts}_{\text{sep}} \rightarrow \mathcal{T}\text{-ReprCat}_{\text{sep}} \rightarrow \mathbf{Set}^{\mathbb{T}},$$

which commute with the canonical forgetful functors to $\mathbf{Set}$. The composite $\mathcal{T}\text{-CoCts}_{\text{sep}} \rightarrow \mathbf{Set}^{\mathbb{T}}$ is even monadic since both $\mathcal{T}\text{-CoCts}_{\text{sep}}$ and $\mathbf{Set}^{\mathbb{T}}$ are monadic over $\mathbf{Set}$ (see [Linton, 1969]).

4. THE VIETORIS MONAD $\mathbb{V}$

General assumption. *From now on until the end of this paper, $\mathcal{T} = (\mathbb{T}, \mathbf{V}, \xi)$ denotes a (strict) topological theory where, moreover, $T1 = 1$.*

Under these conditions, the $\mathcal{T}$ -category $\mathbf{V} = (\mathbf{V}, \text{hom}_\xi) \simeq P1$ is totally cocomplete. Furthermore, since $e_1 : 1 \rightarrow T1$ is a bijection, we can identify a $\mathbf{V}$ -relation $\varphi : 1 \multimap X$ with the $\mathcal{T}$ -relation $\varphi \cdot e_1^\circ : 1 \multimap X$. If, moreover, $X = (X, a)$ is a $\mathcal{T}$ -category, then $\varphi \cdot e_1^\circ$ is a $\mathcal{T}$ -distributor of type $G \multimap X$ if and only if

$$a \cdot T_\xi \varphi \cdot e_1 \leq \varphi.$$

Note that $a \cdot T_\xi \varphi \cdot e_1 \geq \varphi$ holds for every $\mathbf{V}$ -relation $\varphi : 1 \multimap X$.

Lemma 4.1. *For every $\mathcal{T}$ -category $X = (X, a)$, the $\mathcal{T}$ -graph $\mathbf{V}^X$ is dualisable.*

Proof. From Proposition 1.7 we know that the underlying $\mathbf{V}$ -graph structure of $\mathbf{V}^X$ is transitive. Furthermore,

$$\begin{aligned}
\mathbf{V}^X(\mathbf{p}, h) &= \bigwedge_{x \in X} \bigwedge_{\mathfrak{x} \in TX} \bigwedge_{\substack{\mathfrak{w} \in T(\mathbf{V}^X \times X) \\ T\pi_1(\mathfrak{w}) = \mathbf{p}, T\pi_2(\mathfrak{w}) = \mathfrak{x}}} \text{hom}(a(\mathfrak{x}, x), \text{hom}(\xi \cdot T\text{ev}(\mathfrak{w}), h(x))) \\
&= \bigwedge_{x \in X} \bigwedge_{\mathfrak{x} \in TX} \bigwedge_{\substack{\mathfrak{w} \in T(\mathbf{V}^X \times X) \\ T\pi_1(\mathfrak{w}) = \mathbf{p}, T\pi_2(\mathfrak{w}) = \mathfrak{x}}} \text{hom}(a(\mathfrak{x}, x) \otimes \xi \cdot T\text{ev}(\mathfrak{w}), h(x)) \\
&= \bigwedge_{x \in X} \text{hom}\left(\bigvee_{\mathfrak{x} \in TX} a(\mathfrak{x}, x) \otimes \bigvee_{\substack{\mathfrak{w} \in T(\mathbf{V}^X \times X) \\ T\pi_1(\mathfrak{w}) = \mathbf{p}, T\pi_2(\mathfrak{w}) = \mathfrak{x}}} \xi \cdot T\text{ev}(\mathfrak{w}), h(x)\right) \\
&= \bigwedge_{x \in X} \text{hom}(a \cdot T_\xi \text{ev}(\mathbf{p}, x), h(x)) \\
&= [a \cdot T_\xi \text{ev}(\mathbf{p}, -), h],
\end{aligned}$$

where in the last line we consider $\text{ev} : \mathbf{V}^X \times X \rightarrow \mathbf{V}$ as a $\mathbf{V}$ -relation $\text{ev} : \mathbf{V}^X \dashrightarrow X$. Finally, writing $i_{\mathbf{p}} : 1 \rightarrow T(\mathbf{V}^X)$ for the mapping sending the unique point of 1 to $\mathbf{p} \in T(\mathbf{V}^X)$, the composite

$$1 \xrightarrow{i_{\mathbf{p}}} T(\mathbf{V}^X) \xrightarrow{T_\xi \text{ev}} TX \xrightarrow{a} X$$

is a $\mathcal{T}$ -distributor of type $G \dashrightarrow X$ since

$$a \cdot T_\xi a \cdot T_\xi T_\xi \text{ev} \cdot T i_{\mathbf{p}} \cdot e_1 \leq a \cdot T_\xi a \cdot m_X^\circ \cdot T_\xi \text{ev} \cdot i_{\mathbf{p}} = a \cdot T_\xi \text{ev} \cdot i_{\mathbf{p}}.$$

Therefore $\mu(\mathbf{p}) := a \cdot T_\xi \text{ev}(\mathbf{p}, -)$ belongs to $\mathbf{V}^X$. $\square$

Proposition 4.2. *For each core-compact $\mathcal{T}$ -category $X = (X, a)$, $\mathbf{V}^X$ is a separated representable $\mathcal{T}$ -category where the left adjoint of the $\mathcal{T}$ -functor $e_{\mathbf{V}^X} : \mathbf{V}^X \rightarrow T(\mathbf{V}^X)$ is given by*

$$\mu : T(\mathbf{V}^X) \rightarrow \mathbf{V}^X, \mathbf{p} \mapsto a \cdot T_\xi \text{ev}(\mathbf{p}, -).$$

Example 4.3. We consider $\mathcal{T} = \mathcal{U}_2$, and let X be a topological space. We write $\mathcal{O}$ for the collection of all open subsets of X , and $\mathcal{O}(x)$ for the set of all open neighbourhoods of $x \in X$. We can identify 2^X with the set of all closed subsets of X . For any subset $V \subseteq X$, we put

$$V^\diamond = \{A \in 2^X \mid A \cap V \neq \emptyset\}.$$

For an ultrafilter $\mathbf{p}$ on 2^X , the smallest convergence point $\mu(\mathbf{p})$ of $\mathbf{p}$ can be calculated as

$$\begin{aligned}
\mu(\mathbf{p}) &= \{x \in X \mid \exists \mathfrak{x} \in UX. \mathbf{p}(U_\xi \text{ev}) \mathfrak{x} \ \& \ \mathfrak{x} \rightarrow x\} \\
&= \{x \in X \mid \forall V \in \mathcal{O}(x), \mathcal{A} \in \mathbf{p} \exists A \in \mathcal{A}, y \in V. y \in A\} \\
&= \{x \in X \mid \forall V \in \mathcal{O}(x), \mathcal{A} \in \mathbf{p}. V^\diamond \cap \mathcal{A} \neq \emptyset\} \\
&= \{x \in X \mid \forall V \in \mathcal{O}(x). V^\diamond \in \mathbf{p}\}.
\end{aligned}$$

Therefore, for any $A \in (2^X)^{\text{op}}$,

$$\begin{aligned}
\mathbf{p} \rightarrow A &\iff A \subseteq \mu(\mathbf{p}) \\
&\iff \forall x \in A, V \in \mathcal{O}(x). V^\diamond \in \mathbf{p} \\
&\iff \forall V \in \mathcal{O}. (V \cap A \neq \emptyset \Rightarrow V^\diamond \in \mathbf{p}) \\
&\iff \forall V \in \mathcal{O}. (A \in V^\diamond \Rightarrow V^\diamond \in \mathbf{p});
\end{aligned}$$

hence the convergence of the pseudo-topological space $(2^X)^{\text{op}}$ is induced by $\{V^\diamond \mid V \in \mathcal{O}\}$ and therefore $VX := (2^X)^{\text{op}}$ is actually a topological space. This topology on the set of closed subsets of a topological space is known as the *upper Vietoris topology* (see [Clementino and Tholen, 1997], for instance). We find it remarkable that, albeit 2^X belongs to **Top** if and only if X is exponentiable (see [Schwarz, 1984]), its dual $(2^X)^{\text{op}}$ belongs always to **Top**. In fact, we

can now easily derive the well-known characterisation of exponentiable spaces as precisely the core-compact ones (see [Day and Kelly, 1970] and [Isbell, 1975, 1986]):

$$\begin{aligned} X \text{ is exponentiable} &\iff (VX)^{\text{op}} \text{ is topological} \\ &\iff VX \text{ is core-compact} && [\text{Proposition 2.15}] \\ &\iff X \text{ is core-compact.} \end{aligned}$$

The last equivalence follows from the “Sub-Base Lemma” of Example 1.9 (applied to the sub-base $\{V^\diamond \mid V \in \mathcal{O}\}$ of VX).

We also note that $V^\diamond \cap \mathcal{A} \neq \emptyset$ is equivalent to $V \cap \bigcup \mathcal{A} \neq \emptyset$, and therefore

$$\begin{aligned} \mu(\mathfrak{p}) &= \{x \in X \mid \forall \mathcal{A} \in \mathfrak{p}, V \in \mathcal{O}(x). V \cap \bigcup \mathcal{A} \neq \emptyset\} \\ &= \{x \in X \mid \forall \mathcal{A} \in \mathfrak{p}. x \in \overline{\bigcup \mathcal{A}}\} \\ &= \bigcap_{\mathcal{A} \in \mathfrak{p}} \overline{\bigcup \mathcal{A}}. \end{aligned}$$

Here $\overline{(-)}$ denotes the closure of the topological space X . For $K \subseteq X$ compact, $K^\diamond$ is a compact down-set in VX and therefore its complement is open in VX^{op} . Furthermore, for X locally compact, one easily verifies that the sets

$$(K^\diamond)^\complement = \{A \in VX \mid A \cap K = \emptyset\} \quad (K \subseteq X \text{ compact})$$

generate the convergence of VX^{op} (defined by $\mathfrak{p} \rightarrow A \iff \mu(\mathfrak{p}) \subseteq A$), which confirms that the topology of $2^X \simeq (VX)^{\text{op}}$ is the compact-open topology.

Unfortunately, I do not know if $(V^X)^{\text{op}}$ is in general a $\mathcal{T}$ -category. Therefore we will consider mostly core-compact $\mathcal{T}$ -categories in the sequel.

Recall that the structure a of a $\mathcal{T}$ -category $X = (X, a)$ can be seen as a $\mathcal{T}$ -functor $a : (TX)^{\text{op}} \otimes X \rightarrow V$, and therefore, if X is core-compact, induces a $\mathcal{T}$ -functor

$$\ulcorner a \urcorner : (TX)^{\text{op}} \rightarrow V^X.$$

Furthermore, for every $\mathfrak{X} \in TTX$,

$$\ulcorner a \urcorner \cdot m_X(\mathfrak{X}) = a(m_X(\mathfrak{X}), -)$$

and

$$\mu \cdot T \ulcorner a \urcorner (\mathfrak{X}) = a \cdot T_\xi \text{ev}(T \ulcorner a \urcorner_X(\mathfrak{X}), -) = a \cdot T_\xi a(\mathfrak{X}, -),$$

where the last equality follows from the fact that

$$\begin{array}{ccc} TX \times X & \xrightarrow{\ulcorner a \urcorner \times 1_X} & V^X \times X \\ & \searrow a & \downarrow \text{ev} \\ & & V \end{array}$$

commutes. Hence:

Proposition 4.4. *For each core-compact $\mathcal{T}$ -category $X = (X, a)$, $\ulcorner a \urcorner : (TX)^{\text{op}} \rightarrow V^X$ is a morphism in $(V\text{-Cat})^\mathbb{T}$.*

By the result above, $\ulcorner a \urcorner : (TX)^{\text{op}} \rightarrow V^X$ can be dualised to $\Lambda_X := \ulcorner a \urcorner^{\text{op}} : TX \rightarrow (V^X)^{\text{op}}$. But TX is the free $\mathbb{T}$ -algebra on the $\mathcal{T}$ -category $X = (X, a)$, hence Λ_X is completely determined by the *Yoneda map*

$$\hbar_X := \Lambda_X \cdot e_X : X \rightarrow (V^X)^{\text{op}}, \quad x \mapsto a_0(x, -).$$

We put $VX := (V^X)^{\text{op}}$, and denote the structure on VX by $\langle\langle -, - \rangle\rangle$. Hence, for $\mathfrak{p} \in TVX$ and $\varphi \in VX$,

$$\langle\langle \mathfrak{p}, \varphi \rangle\rangle = [\varphi, \mu(\mathfrak{p})].$$

For $\mathfrak{h}_X : X \rightarrow VX$ one has

$$\mathfrak{h}_X^{\otimes}(\mathfrak{p}, x) = \langle\langle \mathfrak{p}, \mathfrak{h}_X(x) \rangle\rangle = [\mathfrak{h}_X(x), \mu(\mathfrak{p})] = \mu(\mathfrak{p})(x), \quad (4)$$

for all $x \in X$ and $\mathfrak{p} \in T(V^X)$; where the last equality follows from the Yoneda Lemma for $\mathbf{V}$ -categories.

Example 4.5. For both $\mathcal{T} = \mathcal{U}_{\mathbf{P}_+}$ and $\mathcal{T} = \mathcal{U}_{\mathbf{P}_\wedge}$ and a core-compact topological space X (viewed as an approach space), the underlying set of the approach space VX is the set of all lower semi-continuous functions from X to $[0, \infty]$ (see [Lowen, 1997, Proposition 2.1.8]).

Lemma 4.6. *For each core-compact $\mathcal{T}$ -category $X = (X, a)$, $\mathfrak{h}_X : X \rightarrow VX$ is fully faithful. If, moreover, X is representable, then $\mathfrak{h}_X : X \rightarrow VX$ is even a pseudo-homomorphism.*

Proof. Since the diagram

$$\begin{array}{ccc} X \times X & \xrightarrow{\mathfrak{h}_X \times 1_X} & \mathbf{V}^X \times X \\ & \searrow a_0 & \downarrow \text{ev} \\ & & \mathbf{V} \end{array}$$

commutes, for $\mathfrak{p} = T \mathfrak{h}_X(\mathfrak{x})$ with $\mathfrak{x} \in TX$ one has (where $x \in X$)

$$\mu(\mathfrak{p})(x) = a \cdot T_\xi \text{ev}(T \mathfrak{h}_X(\mathfrak{x}), x) = a \cdot T_\xi a_0(\mathfrak{x}, x) = a(\mathfrak{x}, x)$$

and therefore $\langle\langle T \mathfrak{h}_X(\mathfrak{x}), \mathfrak{h}_X(x) \rangle\rangle = a(\mathfrak{x}, x)$. To see the second statement, just note that

$$\mu \cdot T \mathfrak{h}_X(\mathfrak{x}) = a(\mathfrak{x}, -) = a_0(\alpha(\mathfrak{x}), -) = \mathfrak{h} \cdot \alpha(\mathfrak{x}),$$

where $\alpha : TX \rightarrow X$ is the $\mathbb{T}$ -algebra structure of X . □

Albeit liftings of $\mathcal{T}$ -distributors do not exist in general in $\mathcal{T}\text{-Dist}$, it is shown in [Hofmann and Waszkiewicz, 2011] that $\mathcal{T}\text{-Dist}$ admits liftings of $\mathcal{T}$ -distributors along $\mathcal{T}$ -distributors of type $1 \multimap X$.

Lemma 4.7. *For all $\mathcal{T}$ -distributors $\varphi : Y \multimap X$ and $\psi : G \multimap X$, φ has a lifting $\psi \multimap \varphi$ along ψ in $\mathcal{T}\text{-Dist}$ which is given by $\psi \multimap \varphi = \psi \cdot e_1 \multimap \bullet \varphi$.*

$$\begin{array}{ccc} X & \xleftarrow{\phi} & Y \\ \psi \downarrow & \swarrow \psi \multimap \phi & \\ G & & \end{array}$$

Every $u \in \mathbf{V}$ can be interpreted as a $\mathcal{T}$ -distributor $u : 1 \multimap 1$, and then $u \otimes v$ corresponds to $v \circ u$. Liftings can be used to turn the ordered set $\mathcal{T}\text{-Dist}(G, X)$ into a $\mathbf{V}$ -category by putting

$$[\varphi, \varphi'] = \varphi \multimap \varphi' = \bigwedge_{x \in X} \text{hom}(\varphi(x), \varphi'(x)),$$

for all $\varphi, \varphi' : G \multimap X$. Hence, for $X = (X, a)$ core-compact, the $\mathbf{V}$ -category $\mathcal{T}\text{-Dist}(G, X)$ is just the dual of the underlying $\mathbf{V}$ -category $(VX)_0$ of VX . For every $\mathcal{T}$ -distributor $\psi : X \multimap Y$, composition with ψ defines a mapping

$$\psi \circ - : \mathcal{T}\text{-Dist}(G, X) \rightarrow \mathcal{T}\text{-Dist}(G, Y)$$

which is actually a $\mathbf{V}$ -functor since

$$\varphi \multimap \varphi' \leq (\psi \circ \varphi) \multimap (\psi \circ \varphi')$$

follows from

$$\psi \circ \varphi \circ (\varphi \multimap \varphi') \leq \psi \circ \varphi'.$$

For a $\mathcal{T}$ -distributor $\psi : X \multimap Y$ between representable $\mathcal{T}$ -categories one might hope that $\psi \circ -$ is a $\mathcal{T}$ -functor of type $VX \rightarrow VY$; unfortunately, this is in general not the case (see Proposition 4.20). Fortunately, the situation is better if $\psi = f^{\otimes}$ for a $\mathcal{T}$ -functor $f : X \rightarrow Y$, as we show next.

Proposition 4.8. *Let $f : X \rightarrow Y$ be a $\mathcal{T}$ -functor between core-compact $\mathcal{T}$ -categories $X = (X, a)$ and $Y = (Y, b)$. Then $Vf := f_{\otimes} \circ - : VX \rightarrow VY$ is a $\mathcal{T}$ -functor. If, moreover, X and Y are representable and f is a pseudo-homomorphism, then Vf is a homomorphism.*

Proof. We put $\Phi := f_{\otimes} \circ -$, and show first that

$$\begin{array}{ccc} V^X & \xrightarrow{\Phi} & V^Y \\ \text{ev}_X \downarrow & \leq & \downarrow \text{ev}_Y \\ X & \xrightarrow{b_0 \cdot f} & Y \end{array}$$

$b_0 \cdot f \cdot \text{ev}_X \leq \text{ev}_Y \cdot \Phi$, with equality if X and Y are representable and f is a pseudo-homomorphism. In fact, for $\varphi \in V^X$ and $y \in Y$,

$$b_0 \cdot f \cdot \text{ev}_X(\varphi, y) = \bigvee_{x \in X} \varphi(x) \otimes b_0(f(x), y)$$

and

$$\text{ev}_Y \cdot \Phi(\varphi, y) = \bigvee_{\mathfrak{x} \in TX} \xi \cdot T\varphi(\mathfrak{x}) \otimes b(Tf(\mathfrak{x}), y).$$

Restricting the second formula to elements of the form $\mathfrak{x} = e_X(x)$ with $x \in X$ gives the first claim. Assume now that X and Y are representable with $a = a_0 \cdot \alpha$ and $b = b_0 \cdot \beta$ and that f is a pseudo-homomorphism. Since $\varphi : X \rightarrow V$ is a $\mathcal{T}$ -functor, $\xi(T\varphi(\mathfrak{x})) \leq \varphi(\alpha(\mathfrak{x}))$ for any $\mathfrak{x} \in TX$ (see Lemma 2.9), and therefore

$$\bigvee_{\mathfrak{x} \in TX} \xi \cdot T\varphi(\mathfrak{x}) \otimes b(Tf(\mathfrak{x}), y) \leq \bigvee_{\mathfrak{x} \in TX} \varphi(\alpha(\mathfrak{x})) \otimes b_0(f(\alpha(\mathfrak{x})), y) = \bigvee_{x \in X} \varphi(x) \otimes b_0(f(x), y).$$

Hence, for any $\mathfrak{p} \in T(V^X)$,

$$\begin{aligned} \Phi \cdot \mu_X(\mathfrak{p}) &= f_{\otimes} \cdot T_{\xi} a \cdot T_{\xi} T_{\xi} \text{ev}_X \cdot T i_{\mathfrak{p}} \cdot e_1 = f_{\otimes} \cdot T_{\xi} a \cdot T_{\xi} T_{\xi} \text{ev}_X \cdot e_{T(V^X)} \cdot i_{\mathfrak{p}} \\ &\leq f_{\otimes} \cdot T_{\xi} a \cdot T_{\xi} T_{\xi} \text{ev}_X \cdot m_{V^X}^{\circ} \cdot i_{\mathfrak{p}} = f_{\otimes} \cdot T_{\xi} a \cdot m_X^{\circ} \cdot T_{\xi} \text{ev}_X \cdot i_{\mathfrak{p}} = f_{\otimes} \cdot T_{\xi} \text{ev}_X \cdot i_{\mathfrak{p}} \end{aligned}$$

and

$$\mu_Y \cdot T\Phi(\mathfrak{p}) = b \cdot T_{\xi} \text{ev}_Y \cdot T\Phi(\mathfrak{p}, -) \geq b \cdot T_{\xi} b_0 \cdot Tf \cdot T_{\xi} \text{ev}_X(\mathfrak{p}, -) = f_{\otimes} \cdot T_{\xi} \text{ev}_X(\mathfrak{p}, -);$$

and we conclude that Φ is indeed a $\mathcal{T}$ -functor of type $(V^X)^{\text{op}} \rightarrow (V^Y)^{\text{op}}$. If $f : X \rightarrow Y$ is a pseudo-homomorphism between representable $\mathcal{T}$ -categories, then the second inequality is actually an equality thanks to the calculations above. Finally,

$$f_{\otimes} \cdot T_{\xi} a \cdot T_{\xi} T_{\xi} \text{ev}_X \cdot T i_{\mathfrak{p}} \cdot e_1 \geq f_{\otimes} \cdot e_X \cdot a \cdot T_{\xi} \text{ev}_X \cdot i_{\mathfrak{p}}$$

and

$$f_{\otimes} \cdot e_X \cdot a = b_0 \cdot f \cdot a \leq b_0 \cdot f \cdot \alpha = b_0 \cdot b \cdot Tf = f_{\otimes};$$

and this shows that also the first inequality becomes also an equality in this case. $\square$

Remark 4.9. If $f : (X, a) \rightarrow (Y, b)$ is a pseudo-homomorphism (where $a = a_0 \cdot \alpha$) and $\varphi : Z \multimap X$, then

$$f_{\otimes} \circ \varphi = b \cdot Tf \cdot T_{\xi} \varphi \cdot m_Z^{\circ} = b_0 \cdot f \cdot \alpha \cdot T_{\xi} \varphi \cdot m_Z^{\circ} = b_0 \cdot f \cdot \varphi = f_{*} \cdot \varphi.$$

Therefore V can be seen as an endofunctor

$$V : \mathcal{T}\text{-Cat}_{\text{core}} \rightarrow \mathcal{T}\text{-Cat}_{\text{core}}$$

on $\mathcal{T}\text{-Cat}_{\text{core}}$, and also as an endofunctor

$$V : \mathcal{T}\text{-ReprCat} \rightarrow \mathcal{T}\text{-ReprCat}$$

on $\mathcal{T}\text{-ReprCat}$. In both cases $\hbar = (\hbar_X)$ is a natural transformation $\hbar : 1 \rightarrow V$ since $f_{\otimes} \circ x_{\otimes} = f(x)_{\otimes}$. We will now show that V is part of a monad on both $\mathcal{T}\text{-ReprCat}$ and $\mathcal{T}\text{-Cat}_{\text{core}}$.

For every core-compact $\mathcal{T}$ -category $X = (X, a)$, the $\mathcal{T}$ -distributor $h_X^\otimes : VX \multimap X$ induces a $\mathbf{V}$ -functor

$$h_X^\otimes \circ - : \mathbf{V}^{(VX)} \rightarrow \mathbf{V}^X$$

which sends $\Phi : VX \rightarrow \mathbf{V}$ to

$$h_X^\otimes \circ \Phi = h_X^\circ \cdot \Phi = \Phi(h_X(-)).$$

Hence, $h_X^\otimes \circ -$ is the mate of the composite

$$\mathbf{V}^{(VX)} \otimes X \xrightarrow{1 \otimes h_X} \mathbf{V}^{(VX)} \otimes VX \xrightarrow{\text{ev}} \mathbf{V}$$

and consequently a $\mathcal{T}$ -functor of type $(VVX)^{\text{op}} \rightarrow (VX)^{\text{op}}$. Note that VX is representable, therefore h_{VX} can be dualised to a $\mathcal{T}$ -functor $h_{VX}^{\text{op}} : (VX)^{\text{op}} \rightarrow (VVX)^{\text{op}}$.

Lemma 4.10. *For every core-compact $\mathcal{T}$ -category $X = (X, a)$, the $\mathcal{T}$ -functor*

$$h_X^\otimes \circ - : (VVX)^{\text{op}} \rightarrow (VX)^{\text{op}}$$

is left adjoint to $h_{VX} : (VX)^{\text{op}} \rightarrow (VVX)^{\text{op}}$.

Proof. Let first $\varphi \in \mathbf{V}^X$. Then

$$h_X^\otimes \circ \varphi_\otimes = [h_X(-), \varphi] = \varphi,$$

hence $(h_X^\otimes \circ -) \cdot e_{VX} = 1$. Let now $\Phi \in \mathbf{V}^{(VX)}$. For every $\varphi \in \mathbf{V}^X$ and $x \in X$,

$$[x_\otimes, \varphi] \otimes \Phi(\varphi) \leq \Phi(x_\otimes)$$

since $\Phi : \mathbf{V}^X \rightarrow \mathbf{V}$ is a $\mathcal{T}$ -functor, hence

$$\Phi(\varphi) \leq \bigwedge_{x \in X} \text{hom}(\varphi(x), \Phi(x_\otimes))$$

for all $\varphi \in \mathbf{V}^X$, that is $\Phi \leq e_{VX} \cdot (h_X^\otimes \circ -)(\Phi)$. $\square$

Corollary 4.11. *For every core-compact $\mathcal{T}$ -category $X = (X, a)$, $h_X^\otimes \circ - : (VVX)^{\text{op}} \rightarrow (VX)^{\text{op}}$ is a homomorphism, and therefore has a dual $\mathcal{T}$ -functor $\omega_X := (h_X^\otimes \circ -)^{\text{op}} : VVX \rightarrow VX$. Furthermore,*

$$h_{VX} \dashv \omega_X \dashv V h_X$$

in $\mathcal{T}\text{-Cat}$ and, if X is representable, even in $\mathcal{T}\text{-ReprCat}$.

Proposition 4.12. $\omega = (\omega_X) : VV \rightarrow V$ is a natural transformation where $V : \mathcal{T}\text{-ReprCat} \rightarrow \mathcal{T}\text{-ReprCat}$.

Proof. Let $f : X \rightarrow Y$ in $\mathcal{T}\text{-ReprCat}$. We show first that $h_Y^\otimes \circ (Vf)_\otimes = f_\otimes \circ h_X^\otimes$. Let $\mathbf{p} \in T(\mathbf{V}^X)$ and $y \in Y$. Then

$$\begin{aligned} h_Y^\otimes \circ (Vf)_\otimes(\mathbf{p}, y) &= h_Y^\otimes(TVf(\mathbf{p}), y) \\ &= \mu_Y(TVf(\mathbf{p}))(y) && \text{by (4)} \\ &= Vf(\mu_X(\mathbf{p}))(y) && Vf \text{ is a homomorphism} \\ &= f_\otimes \circ \mu_X(\mathbf{p})(y) && \text{definition of } Vf \\ &= f_* \cdot \mu_X(\mathbf{p})(y) && \text{Remark 4.9} \\ &= f_* \cdot h_X^\otimes(\mathbf{p}, y) && \text{by (4)} \\ &= f_\otimes \circ h_X^\otimes(\mathbf{p}, y). \end{aligned}$$

Hence, for all $\Phi \in VVX$,

$$Vf(\omega_X(\Phi)) = f_\otimes \circ h_X^\otimes \circ \Phi = h_Y^\otimes \circ (Vf)_\otimes \circ \Phi = \omega_Y(VVf(\Phi)). \quad \square$$

All told:

Theorem 4.13. $\mathbb{V} = (V, h, \omega)$ is a monad on $\mathcal{T}\text{-ReprCat}$ which, moreover, is of Kock-Zöberlein type.

Remark 4.14. The monad $\mathbb{V} = (V, h, \omega)$ on $\mathcal{T}\text{-ReprCat}$ restricts to a monad $\mathbb{V} = (V, h, \omega)$ on $\mathcal{T}\text{-ReprCat}_{\text{sep}}$ since VX is separated, for every $\mathcal{T}$ -category X ; and the categories $(\mathcal{T}\text{-ReprCat})^{\mathbb{V}}$ and $(\mathcal{T}\text{-ReprCat}_{\text{sep}})^{\mathbb{V}}$ of Eilenberg–Moore algebras are actually equal. Furthermore, we can lift $\mathbb{V}$ to a monad $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on $\text{Set}^{\mathbb{T}}$ via the adjunction (see Remark 2.7)

$$\mathcal{T}\text{-ReprCat}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad \quad} \end{array} \text{Set}^{\mathbb{T}},$$

that is, $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ is the monad on $\text{Set}^{\mathbb{T}}$ induced by the composite of the adjunctions

$$(\mathcal{T}\text{-ReprCat}_{\text{sep}})^{\mathbb{V}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad \quad} \end{array} \mathcal{T}\text{-ReprCat}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad \quad} \end{array} \text{Set}^{\mathbb{T}}.$$

Explicitly, the unit $\tilde{h}_X : X \rightarrow \tilde{V}X$ at X is defined by $i \cdot \tilde{h}_X = h_X$, where $i : \tilde{V}X \rightarrow VX$, $\varphi \mapsto \varphi$; and the multiplication $\tilde{\omega}_X$ at X sends $\Phi : G \rightarrow \tilde{V}X$ to $h_X^{\circ} \circ i_{\circ} \circ \Phi$. We shall see in Remark 7.5 that $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ is also induced by the adjunction

$$\mathcal{T}\text{-CoCts}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad \quad} \end{array} \text{Set}^{\mathbb{T}}$$

of Remark 3.9.

Example 4.15. In the topological case, the monad $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on CompHaus is the Vietoris monad whose functor part was originally studied in Vietoris [1922]. In the approach case (i.e. $\mathcal{T} = \mathcal{U}_{\mathcal{P}_+}$), we obtain a monad $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on CompHaus where $\tilde{V}X$ is the compact Hausdorff space with underlying set all lower semi-continuous functions of type $X \rightarrow [0, \infty]$ (see Example 4.5), and where an ultrafilter $\mathfrak{p}$ on $\tilde{V}X$ converges to (see Proposition 4.2)

$$X \rightarrow [0, \infty], x \mapsto \inf \left\{ \sup_{\mathcal{A} \in \mathfrak{p}, A \in \mathfrak{t}} \inf_{\varphi \in \mathcal{A}, z \in A} \varphi(z) \mid \mathfrak{t} \in UX, \mathfrak{t} \rightarrow x \right\}.$$

The monad $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on CompHaus obtained from $\mathcal{U}_{\mathcal{P}_\Lambda}$ has the same functor and the same unit as for $\mathcal{U}_{\mathcal{P}_+}$, but the multiplication is different.

Comparing with the situation for $\mathbf{V}$ -categories, one might expect $Vf : VX \rightarrow VY$ to be right adjoint. If it is so, its left adjoint is necessarily given by $f^{\circ} \circ -$, which is the dual of the exponential $V^f : V^Y \rightarrow V^X$, for $f : X \rightarrow Y$. Therefore we have to investigate whether or not $Vf : V^Y \rightarrow V^X$ is a homomorphism; as it turns out, this is only true under additional conditions on f .

Definition 4.16. A $\mathcal{T}$ -functor $f : (X, a) \rightarrow (Y, b)$ is called *downwards open* whenever $a \cdot Tf^{\circ} \cdot T_{\xi}b_0 \geq f^{\circ} \cdot b$.

Note that every $\mathcal{T}$ -functor $f : (X, a) \rightarrow (Y, b)$ satisfies $a \cdot Tf^{\circ} \cdot T_{\xi}b_0 \leq f^{\circ} \cdot b$, hence for a downwards open $\mathcal{T}$ -functor one actually has equality. Of course, every open $\mathcal{T}$ -functor (see [Clementino and Hofmann, 2004, Definition 4.1]) is downwards open, and the reverse is true whenever $b_0 = 1_Y$. In particular, any non-open continuous map between compact Hausdorff spaces is an example of a homomorphism of representable spaces which is not downwards open.

Lemma 4.17. *For every $\mathbf{V}$ -functor $f : (X, c) \rightarrow (Y, d)$, the $\mathcal{T}$ -functor $f : (X, e_X^{\circ} \cdot T_{\xi}c) \rightarrow (Y, e_Y^{\circ} \cdot T_{\xi}d)$ is downwards open. In particular, every $\mathbf{V}$ -functor is downwards open in $\mathbf{V}\text{-Cat}$.*

Proof. We put $a = e_X^{\circ} \cdot T_{\xi}c$ and $b = e_Y^{\circ} \cdot T_{\xi}d$. Then $f^{\circ} \cdot b = e_X^{\circ} \cdot Tf^{\circ} \cdot T_{\xi}d \leq e_X^{\circ} \cdot Tf^{\circ} \cdot T_{\xi}b_0$. $\square$

Lemma 4.18. *For every $\mathcal{T}$ -category $X = (X, a)$, $e_X : X \rightarrow TX$ is downwards open if and only if X is core-compact.*

Proof. Recall that the $\mathcal{T}$ -category structure on TX is given by $T_{\xi}a \cdot m_X^{\circ} \cdot m_X$, and the underlying $\mathbf{V}$ -category structure is $T_{\xi}a \cdot m_X^{\circ}$. We compute

$$e_X^{\circ} \cdot T_{\xi}a \cdot m_X^{\circ} \cdot m_X = (e_X \circ a) \cdot m_X = a \cdot m_X$$

and

$$a \cdot Te_X^\circ \cdot T_\xi(T_\xi a \cdot m_X^\circ) = a \cdot T_\xi(e_X \circ a) = a \cdot T_\xi a,$$

which proves the claim. $\square$

Example 4.19. A continuous map $f : X \rightarrow Y$ between topological spaces is downwards open if and only if the down closure $\downarrow f(A)$ of every open subset $A \subseteq X$ is open in Y . To see this, we recall (Example 2.1) that

$$\mathfrak{x}(U_\xi \leq) \mathfrak{x}' \iff \forall A \in \mathfrak{x}'. \downarrow A \in \mathfrak{x},$$

where $\leq$ is an order relation on X and $\mathfrak{x}, \mathfrak{x}' \in UX$. Assume first that f is downwards open and let $A \subseteq X$ open. Let $\mathfrak{y} \rightarrow y \leq f(x)$ in Y , for some $x \in A$. By hypothesis, there is some $\mathfrak{x} \in UX$ with $\mathfrak{x} \rightarrow x$ and $\mathfrak{y}(U_\xi \leq) Uf(\mathfrak{x})$. From A open it follows that $A \in \mathfrak{x}$, hence $\downarrow f(A) \in \mathfrak{y}$. Conversely, assume now that $\downarrow f(A)$ is open, for every open subset $A \subseteq X$. Let $x \in X$ and $\mathfrak{y} \in UY$ with $\mathfrak{y} \rightarrow f(x)$. Then the ideal

$$\{A \subseteq X \mid \downarrow f(A) \notin \mathfrak{y}\}$$

is disjoint from the neighbourhood filter of x , and therefore (see Proposition 1.5) there is some ultrafilter $\mathfrak{x} \in UX$ with $\mathfrak{x} \rightarrow x$ and $\mathfrak{y}(U_\xi \leq) Uf(\mathfrak{x})$.

Proposition 4.20. *Let $f : (X, a) \rightarrow (Y, b)$ be a $\mathcal{T}$ -functor between core-compact $\mathcal{T}$ -categories. Then the following assertions are equivalent.*

- (i) f is downwards open.
- (ii) $V^f : V^Y \rightarrow V^X$ is a homomorphism.
- (iii) $Vf : VX \rightarrow VY$ has a left adjoint.

Proof. Assume first that f is downwards open. Let $\mathfrak{q} \in T(V^Y)$ and $x \in X$. Then

$$V^f(\mu_Y(\mathfrak{q}))(x) = \mu_Y(\mathfrak{q})(f(x)) = b \cdot T_\xi \text{ev}_Y(\mathfrak{q}, f(x)) = f^\circ \cdot b \cdot T_\xi \text{ev}_Y(\mathfrak{q}, x)$$

and

$$\mu_X(T(V^f)(\mathfrak{q}))(x) = a \cdot T_\xi \text{ev}_X(T(V^f)(\mathfrak{q}), x).$$

Furthermore, the diagram

$$\begin{array}{ccc} V^Y & \xrightarrow{V^f} & V^X \\ \text{ev}_Y \downarrow & & \downarrow \text{ev}_X \\ Y & \xrightarrow{f^\circ \cdot b_0} & X \end{array}$$

commutes since, for every $\varphi : V^Y$ and $x \in X$,

$$\text{ev}_X \cdot V^f(\varphi, x) = \varphi(f(x)) = b_0 \cdot \varphi(f(x)) = f^\circ \cdot b_0 \cdot \text{ev}_Y(\varphi, x).$$

Therefore $a \cdot T_\xi \text{ev}_X(T(V^f)(\mathfrak{q}), x) = a \cdot T f^\circ \cdot T_\xi b_0 \cdot T_\xi \text{ev}_Y(\mathfrak{q}, x)$. We conclude that V^f is a homomorphism, hence (i) $\Rightarrow$ (ii). Since $\text{ev}_Y \cdot h_Y = b_0$, from $f^\circ \cdot b \cdot T_\xi \text{ev}_Y = a \cdot T f^\circ \cdot T_\xi b_0 \cdot T_\xi \text{ev}_Y$ one obtains $f^\circ \cdot b = a \cdot T f^\circ \cdot T_\xi b_0$; hence (ii) $\Rightarrow$ (i). Finally, the equivalence (ii) $\Leftrightarrow$ (iii) is clear. $\square$

Corollary 4.21. *Let $f : (X, a) \rightarrow (Y, b)$ be a downwards open $\mathcal{T}$ -functor between core-compact $\mathcal{T}$ -categories. Then the diagram*

$$\begin{array}{ccc} VVX & \xrightarrow{VVf} & VVY \\ \omega_X \downarrow & & \downarrow \omega_Y \\ VX & \xrightarrow{Vf} & VY \end{array}$$

commutes.

Proof. Just note that the diagram of the corresponding left adjoints commutes since $h : 1 \rightarrow V$ is a natural transformation. $\square$

Corollary 4.22. $\omega = (\omega_X) : VV \rightarrow V$ is a natural transformation where $V : \mathcal{T}\text{-Cat}_{\text{core}} \rightarrow \mathcal{T}\text{-Cat}_{\text{core}}$.

Proof. Let $f : (X, a) \rightarrow (Y, b)$ be a $\mathcal{T}$ -functor between $\mathcal{T}$ -categories satisfying $a \cdot T_\xi a = a \cdot m_X$ and $b \cdot T_\xi b = b \cdot m_Y$. Put $g := Tf \cdot e_X = e_Y \cdot f$. Then the naturality square of ω for g commutes since g is the composite of a pseudo-homomorphism and a downwards open $\mathcal{T}$ -functor. Furthermore, $V(e_Y)$ is injective since e_Y is fully faithful, and the naturality square of ω for e_Y commutes since e_Y is downwards open. Consequently, also

$$\begin{array}{ccc} VVX & \xrightarrow{VVf} & VVY \\ \omega_X \downarrow & & \downarrow \omega_Y \\ VX & \xrightarrow{Vf} & VY \end{array}$$

commutes. □

Theorem 4.23. $\mathbb{V} = (V, h, \omega)$ is a Kock-Zöberlein monad on $\mathcal{T}\text{-Cat}_{\text{core}}$.

5. COMPLETE $\mathcal{T}$ -CATEGORIES

Similar to what was done for colimits, we introduce now the notion of weighted limit following closely the $\mathbf{V}$ -categorical case. A *weighted limit diagram* (h, φ) in a $\mathcal{T}$ -category X is given by a $\mathcal{T}$ -functor $h : A \rightarrow X$ and a $\mathcal{T}$ -distributor $\varphi : G \multimap A$,

$$\begin{array}{ccc} A & \xrightarrow{h} & X \\ \varphi \uparrow \circ & & \\ G & & \end{array}$$

and $x_0 \in X$ is a *limit* of this diagram, written as $x_0 \simeq \lim(h, \varphi)$, if x_0 represents $\varphi \multimap h^*$ in the sense that $x_0^* = \varphi \multimap h^*$. We hasten to remark that *we cannot consider an arbitrary $\mathcal{T}$ -distributor $\varphi : B \multimap A$ above since the lifting $\varphi \multimap h^*$ might not exist*. A $\mathcal{T}$ -functor $f : X \rightarrow Y$ *preserves the limit* of h and φ whenever $f(x_0) \simeq \lim(f \cdot h, \varphi)$, and $f : X \rightarrow Y$ is said to be *continuous* whenever f preserves all weighted limits which exist in X . Note that, for any $\mathfrak{x} \in TX$,

$$\varphi \multimap h^*(\mathfrak{x}) = \bigwedge_{z \in A} \text{hom}(\varphi(z), a(\mathfrak{x}, h(z))),$$

hence $x_0 \simeq \lim(h, \varphi)$ precisely when, for all $\mathfrak{x} \in TX$,

$$a(\mathfrak{x}, x_0) = \bigwedge_{z \in A} \text{hom}(\varphi(z), a(\mathfrak{x}, h(z))).$$

In particular, the equality above holds for all $\mathfrak{x} = e_X(x)$, $x \in X$, therefore x_0 is also a limit of the underlying diagram in $\mathbf{V}\text{-Cat}$; however, a $\mathbf{V}$ -categorical limit in X_0 does not need to be a limit in the $\mathcal{T}$ -category X (see Example 5.2). Nevertheless, if we know that a diagram has a limit in X , then this limit can be calculated in the underlying $\mathbf{V}$ -category X_0 .

Remark 5.1. A particular instance of a weighted limit in a topological space was considered in [Lucyshyn-Wright, 2011] and called directed conjunction there.

Example 5.2. We consider the ordered set $X_0 = [0, 1]$ (closed unit interval) with the usual order $\leq$, and let X be the induced Alexandroff space. Hence, the closed subsets of X are precisely the up-closed subsets of $[0, 1]$, and an ultrafilter $\mathfrak{x} \in UX$ converges to $x \in X$ if and only if, for all $A \in \mathfrak{x}$, there exists some $y \in A$ with $y \leq x$. Clearly, 0 is the infimum of $A =]0, 1]$ in X_0 , but we shall see that 0 is not an infimum of the closed subset $A =]0, 1]$ of X in $\mathbf{Top}$. In fact, let $\mathfrak{x}$ be any ultrafilter containing the sets $]0, r]$, for $r > 0$. By construction, $\mathfrak{x}$ converges to every $x \in X$ different from 0, but not to $x = 0$.

A $\mathcal{T}$ -category X is called *complete* whenever every weighted limit diagram in X has a limit. To check for completeness it is enough to consider weighted limit diagrams where h is the identity $1_X : X \rightarrow X$ on X since a limit of (h, φ) is a limit of $(1_X, h^{\otimes} \circ \varphi)$, and vice versa. A limit of $(1_X, \varphi)$ we also call *infimum* of φ and we denote such a limit as $\text{Inf}_X(\varphi)$ (or simply $\text{Inf}(\varphi)$). Note that, for a $\mathcal{T}$ -category $X = (X, a)$ and $\varphi : G \multimap X$, $x_0 \simeq \text{Inf}_X(\varphi)$ precisely when, for all $\mathfrak{x} \in TX$,

$$a(\mathfrak{x}, x_0) = \bigwedge_{x \in X} \text{hom}(\varphi(x), a(\mathfrak{x}, x)).$$

If $f : X \rightarrow Y$ is a $\mathcal{T}$ -functor between complete $\mathcal{T}$ -categories, then f is continuous if and only if f sends, for any $\varphi : G \multimap X$, a infimum of φ to a limit of the diagram (f, φ) in Y . Since both limits can be calculated in the underlying $\mathbf{V}$ -categories, we find that the $\mathcal{T}$ -functor f is continuous if the underlying $\mathbf{V}$ -functor is continuous. In fact:

Lemma 5.3. *For every complete $\mathcal{T}$ -category X , the $\mathbf{V}$ -category X_0 is complete. Moreover, a $\mathcal{T}$ -functor $f : X \rightarrow Y$ between complete $\mathcal{T}$ -categories is continuous if and only if the underlying $\mathbf{V}$ -functor $f : X_0 \rightarrow Y_0$ is continuous.*

Proof. Every $\mathbf{V}$ -distributor $\varphi : G \multimap X_0$ can be seen as a $\mathcal{T}$ -distributor $\varphi : G \multimap A(X_0)$, and a limit of the diagram given by $\varphi : G \multimap A(X_0)$, $A(X_0) \rightarrow X$ is also a infimum of $\varphi : G \multimap X_0$ in X_0 . The second claim is now clear. $\square$

The following lemma is dual to [Hofmann, 2011, Lemma 1.8].

Lemma 5.4. *Let $\varphi : G \multimap X$, $\beta : Y \multimap X$ and $\alpha : Z \rightarrow X$ be $\mathcal{T}$ -distributors where α is left adjoint. Then*

$$\varphi \multimap (\beta \circ \alpha) = (\varphi \multimap \beta) \circ \alpha.$$

As expected:

Proposition 5.5. *Every right adjoint $\mathcal{T}$ -functor is continuous.*

Proof. For a right adjoint $\mathcal{T}$ -functor $f : X \rightarrow Y$, a $\mathcal{T}$ -distributor $\varphi : G \multimap A$, a $\mathcal{T}$ -functor $h : A \rightarrow X$ and $x_0 \in X$ with $x_0 \simeq \lim(h, \varphi)$, we calculate

$$\varphi \multimap (f \cdot h)^{\otimes} = (\varphi \multimap h^{\otimes}) \circ f^{\otimes} = x_0^{\otimes} \circ f^{\otimes} = f(x_0)^{\otimes}.$$

Here we use Lemma 5.4 and that $f^{\otimes}$ is a left adjoint $\mathcal{T}$ -distributor since f is right adjoint. $\square$

The following proposition is in sharp contrast to the case of weighted colimits where the existence of all colimits with weights $X \multimap G$ does not guarantee the existence of a left adjoint of the Yoneda embedding $y_X : X \rightarrow PX$ in $\mathcal{T}\text{-Cat}$.

Proposition 5.6. *Let $X = (X, a)$ be a core-compact $\mathcal{T}$ -category. Then X is complete if and only if $\mathfrak{h}_X : X \rightarrow VX$ has a right adjoint $\text{Inf}_X : VX \rightarrow X$ in $\mathcal{T}\text{-Cat}$.*

Proof. First note that

$$\mu_X \cdot T \mathfrak{h}_X = \mu_X \cdot T \Lambda_X \cdot e_X = \Lambda_X \cdot m_X \cdot e_X = \Lambda_X,$$

that is, $a(\mathfrak{x}, -) = \mu_X(T \mathfrak{h}_X(\mathfrak{x}))$ for all $\mathfrak{x} \in TX$. Therefore X is complete if and only if there exists a map $\text{Inf}_X : VX \rightarrow X$ satisfying

$$a(\mathfrak{x}, \text{Inf}_X(\varphi)) = [\varphi, a(\mathfrak{x}, -)] = [\varphi, \mu_X(T \mathfrak{h}_X(\mathfrak{x}))] = \langle T \mathfrak{h}_X(\mathfrak{x}), \varphi \rangle,$$

for all $\mathfrak{x} \in TX$ and $\varphi : G \multimap X$. But this conditions just means that $(\mathfrak{h}_X)_{\otimes} = \text{Inf}_X^{\otimes}$, and the assertion follows using (1) of Subsection VII of Section 1. $\square$

Proposition 5.7. *Let $X = (X, a)$ and $Y = (Y, b)$ be complete and core-compact $\mathcal{T}$ -categories. Then a $\mathcal{T}$ -functor $f : X \rightarrow Y$ is continuous if and only if the diagram*

$$\begin{array}{ccc} VX & \xrightarrow{Vf} & VY \\ \text{Inf}_X \downarrow & \simeq & \downarrow \text{Inf}_Y \\ X & \xrightarrow{f} & Y \end{array}$$

commutes up to equivalence.

Theorem 5.8. *The category $(\mathcal{T}\text{-Cat}_{\text{core}})^{\mathbb{V}}$ of Eilenberg–Moore algebras of $\mathbb{V}$ is precisely the category of core-compact, complete and separated $\mathcal{T}$ -categories and continuous $\mathcal{T}$ -functors.*

Proposition 5.9. *Let $X = (X, a)$ be a core-compact $\mathcal{T}$ -category. Then X is complete if and only if X is representable and X_0 is a complete $\mathbb{V}$ -category.*

Proof. Assume first that X is complete. Then X_0 is complete by Lemma 5.3, and a left inverse of $e_X : X \rightarrow TX$ is given by $\text{Inf}_X \cdot \Lambda_X$. Assume now that X is representable (with algebra structure $\alpha : TX \rightarrow X$) and that X_0 is complete. Let $\varphi : G \multimap X$ be a $\mathcal{T}$ -distributor, then φ can be also seen as a $\mathbb{V}$ -distributor $\varphi : G \multimap X_0$. Let x be an infimum of $\varphi : G \multimap X_0$ in X_0 . Then, for every $\mathfrak{x} \in TX$,

$$a(\mathfrak{x}, x) = a_0(\alpha(\mathfrak{x}), x) = \bigwedge_{x \in X} \text{hom}(\varphi(x), a_0(\alpha(\mathfrak{x}), x)) = \bigwedge_{x \in X} \text{hom}(\varphi(x), a(\mathfrak{x}, x)),$$

hence x is also an infimum of $\varphi : 1 \multimap X$. □

Proposition 5.10. *Let $X = (X, a)$ and $Y = (Y, b)$ be core-compact $\mathcal{T}$ -categories where X is complete. Then the following assertions are equivalent for a $\mathcal{T}$ -functor $f : X \rightarrow Y$.*

- (i) *f is right adjoint.*
- (ii) *f preserves limits and Vf is right adjoint.*
- (iii) *f preserves limits and is downwards open.*

Proof. The equivalence (ii) $\Leftrightarrow$ (iii) follows from Proposition 4.20. The assertion (ii) follows from (i) by Proposition 5.5 and the fact that V is a 2-functor. Finally, assuming (ii), a left adjoint of f is given by $\text{Inf}_X \cdot G \cdot h_Y$ where G is a left adjoint of Vf . □

Corollary 5.11. *Let $X = (X, a)$ be a core-compact $\mathcal{T}$ -category. Then X is complete if and only if X is injective in $\mathcal{T}\text{-Cat}_{\text{core}}$ with respect to fully faithful downwards open $\mathcal{T}$ -functors.*

Proof. This is an instance of [Escardó, 1998, Theorem 4.2.2]. □

6. ISBELL CONJUGATION ADJUNCTION

For every $\mathcal{T}$ -category $X = (X, a)$, there is an adjunction

$$\mathcal{T}\text{-Dist}(G, X)^{\text{op}} \begin{array}{c} \xrightarrow{(-)^-} \\ \Upsilon \\ \xleftarrow{(-)^+} \end{array} \mathcal{T}\text{-Dist}(X, G)$$

in $\mathbb{V}\text{-Cat}$ where

$$\varphi^-(\mathfrak{x}) = \varphi \multimap 1_X^{\otimes}(\mathfrak{x}) = \bigwedge_{x \in X} \text{hom}(\varphi(x), a(\mathfrak{x}, x))$$

for all $\varphi : G \multimap X$ and $\mathfrak{x} \in TX$, and

$$\psi^+(x) = 1_X^{\otimes} \multimap \psi(x) = \bigwedge_{\mathfrak{x} \in TX} \text{hom}(\psi(\mathfrak{x}), a(\mathfrak{x}, x))$$

for all $\psi : X \multimap G$ and $x \in X$. Following [Wood, 2004], we refer to this adjunction as an *Isbell conjugation adjunction*. Note that

$$(x_{\otimes})^- = x^{\otimes} \quad \text{and} \quad (x^{\otimes})^+ = x_{\otimes},$$

for all $x \in X$. Furthermore $x \in X$ is an infimum of $\varphi : G \multimap X$ if and only if $x^{\otimes} = \varphi^-$, and x is a supremum of $\psi : X \multimap G$ if and only if $x_{\otimes} = \psi^+$.

Proposition 6.1. *For every $\mathcal{T}$ -distributor $\varphi : G \multimap X$ and $x \in X$, x is an infimum of φ in X if and only if x is a supremum of φ^- in X . Similarly, for every $\mathcal{T}$ -distributor $\psi : X \multimap G$ and $x \in X$, x is a supremum of ψ if and only if x is an infimum of ψ^+ .*

Proof. If $x^{\otimes} = \varphi^-$, then $x_{\otimes} = (\varphi^-)^+$, and this in turn implies $x^{\otimes} = ((\varphi^-)^+)^- = \varphi^-$. A similar argumentation proves the second claim. $\square$

Theorem 6.2. *A $\mathcal{T}$ -category X is complete if and only if X is cocomplete.*

Lemma 6.3. *Let $f : (X, a) \rightarrow (Y, b)$ be a $\mathcal{T}$ -functor where (Y, b) is representable with $\mathbb{T}$ -algebra structure $\beta : TY \rightarrow Y$. Then*

$$f_{\otimes} = \bar{f}_*,$$

where $\bar{f} : TX \rightarrow Y$ is the extension of $f : (X, a) \rightarrow (Y, b)$ along $e_X : X \rightarrow TX$.

Proof. We calculate:

$$f_{\otimes} = b \cdot Tf = b_0 \cdot \beta \cdot Tf = b_0 \cdot \bar{f} = \bar{f}_*. \quad \square$$

Proposition 6.4. *For every core-compact $\mathcal{T}$ -category $X = (X, a)$, the $\mathbf{V}$ -functor $(-)^-$ is actually a $\mathcal{T}$ -functor $(-)^- : VX \rightarrow PX$.*

Proof. Note that

$$(\lambda_X)_{\otimes}(\mathfrak{x}, \varphi) = (\Lambda_X)_*(\mathfrak{x}, \varphi) = [\varphi, \Lambda_X(\mathfrak{x})] = \bigwedge_{x \in X} \text{hom}(\varphi(x), a(\mathfrak{x}, x)),$$

for all $\varphi \in VX$ and $\mathfrak{x} \in TX$; hence $(-)^-$ is the mate of $(\lambda_X)_{\otimes} : X \multimap VX$. $\square$

However, the $\mathbf{V}$ -functor $(-)^+$ is in general not a $\mathcal{T}$ -functor of type $PX \rightarrow VX$. For instance, the representable approach space $\mathbf{P}_+^{\text{op}}$ is complete (and cocomplete) but not totally cocomplete, and therefore $(-)^+$ cannot be a $\mathcal{T}$ -functor for $X = \mathbf{P}_+^{\text{op}}$.

7. TOTALLY COMPLETE $\mathcal{T}$ -CATEGORIES

At the beginning of Section 5 we pointed already out that the notion of complete $\mathcal{T}$ -category cannot be strengthened to “totally complete” exactly the same was as it was done for cocompleteness, namely by allowing all $\mathcal{T}$ -distributors $\psi : B \multimap A$ as limit weights. Nevertheless, in this section we introduce a notion of total completeness which turns out to be the dual of total cocompleteness.

Definition 7.1. A representable $\mathcal{T}$ -category $X = (X, a)$ is called *totally complete* if $\mathfrak{h}_X : X \rightarrow VX$ has a right adjoint $\text{Inf}_X : VX \rightarrow X$ in $\mathcal{T}\text{-ReprCat}$.

Hence, a totally complete $\mathcal{T}$ -category X is a complete representable $\mathcal{T}$ -category where, moreover, $\text{Inf}_X : VX \rightarrow X$ is a pseudo-homomorphism. We write

$$\mathcal{T}\text{-Cts}$$

for the category of totally complete $\mathcal{T}$ -categories and pseudo-homomorphisms which preserve limits, and $\mathcal{T}\text{-Cts}_{\text{sep}}$ denotes its full subcategory defined by the separated $\mathcal{T}$ -categories. By definition, $\mathcal{T}\text{-Cts}_{\text{sep}} \simeq (\mathcal{T}\text{-ReprCat})^{\mathbf{V}}$. Clearly, every $\mathcal{T}$ -category of the form VX is totally complete. Since $(PX)^{\text{op}} = \mathbf{V}((TX)^{\text{op}})$, this includes the duals of $\mathcal{T}$ -categories of the form PX . In fact, we will show that the totally complete $\mathcal{T}$ -categories are precisely the duals of totally cocomplete $\mathcal{T}$ -categories.

Lemma 7.2. *For every $\mathcal{T}$ -category X , the diagrams of $\mathbf{V}$ -functors*

$$\begin{array}{ccc} X^{\text{op}} & \xrightarrow{e_X^{\text{op}}} & (TX)^{\text{op}} \\ & \searrow y_X^{\text{op}} & \downarrow h_{(TX)^{\text{op}}} \\ & & V((TX)^{\text{op}}) \end{array} \quad \text{and} \quad \begin{array}{ccc} X^{\text{op}} & \xrightarrow{h_{X^{\text{op}}}} & V(X^{\text{op}}) \\ & \searrow y_X^{\text{op}} & \downarrow (\alpha^{\text{op}})^{\otimes} \circ - \\ & & V((TX)^{\text{op}}) \end{array}$$

commute, where in the latter diagram we assume that X is representable with left adjoint $\alpha : TX \rightarrow X$ of $e_X : X \rightarrow TX$.

Proof. Let $X = (X, a)$ be a $\mathcal{T}$ -category, and put $\hat{a} = T_\xi a \cdot m_X^\circ$. Then, for every $x \in X$,

$$h_{(TX)^{\text{op}}}(e_X^{\text{op}}(x)) = \hat{a}^\circ(e_X^{\text{op}}(x), -) = a(-, x) = y_X^{\text{op}}(x) = a_0^\circ(x, \alpha(-)) = (\alpha^{\text{op}})^{\otimes} \circ h_{X^{\text{op}}}(x). \quad \square$$

Proposition 7.3. *Let X be a representable $\mathcal{T}$ -category. Then X is totally cocomplete if and only if X^{op} is totally complete.*

Proof. Assume first that X is totally cocomplete. By Theorem 6.2, X^{op} is complete, we write $\text{Inf}_{X^{\text{op}}} : V(X^{\text{op}}) \rightarrow X^{\text{op}}$ for a right adjoint of $h_{X^{\text{op}}} : X^{\text{op}} \rightarrow V(X^{\text{op}})$ in $\mathcal{T}\text{-Cat}$. We need to prove that $\text{Inf}_{X^{\text{op}}}$ is a pseudo-homomorphism. Firstly, the diagram

$$\begin{array}{ccc} (PX)^{\text{op}} = V((TX)^{\text{op}}) & \xrightarrow{V(\alpha^{\text{op}})} & V(X^{\text{op}}) \\ & \searrow \text{Sup}_X^{\text{op}} & \downarrow \text{Inf}_{X^{\text{op}}} \\ & & X^{\text{op}} \end{array} \quad (5)$$

of $\mathcal{T}$ -functors commutes up to equivalence since the underlying diagram in $\mathbf{V}\text{-Cat}$ consists precisely of the right adjoints of the $\mathbf{V}$ -functors of the second diagram in Lemma 7.2. Since $\text{Sup}_X : PX \rightarrow X$ and $\alpha : TX \rightarrow X$ are left adjoints in $\mathcal{T}\text{-Cat}$, they are in particular pseudo-homomorphisms, hence Sup_X^{op} and $V(\alpha^{\text{op}})$ are pseudo-homomorphisms. Since $V(\alpha^{\text{op}})$ is a split epimorphism in $\mathbf{V}\text{-Cat}$, also $\text{Inf}_{X^{\text{op}}}$ is a pseudo-homomorphism. Conversely, assume now that X^{op} is totally complete. Hence $\text{Inf}_{X^{\text{op}}} : V(X^{\text{op}}) \rightarrow X^{\text{op}}$ is a pseudo-homomorphism. We show that $\text{Inf}_{X^{\text{op}}}^{\text{op}} \cdot V(\alpha^{\text{op}})^{\text{op}}$ is a left adjoint (=left inverse in this case) of $y_X : X \rightarrow PX$. In fact, for the duals of the underlying $\mathbf{V}$ -functors one verifies:

$$\text{Inf}_{X^{\text{op}}} \cdot V(\alpha^{\text{op}}) \cdot y_X^{\text{op}} = \text{Inf}_{X^{\text{op}}} \cdot V(\alpha^{\text{op}}) \cdot h_{(TX)^{\text{op}}} \cdot e_X^{\text{op}} = \text{Inf}_{X^{\text{op}}} \cdot h_{X^{\text{op}}} \cdot \alpha^{\text{op}} \cdot e_X^{\text{op}} \simeq 1_{X_0}. \quad \square$$

From commutativity of (5) we also deduce that a pseudo-homomorphism $f : X \rightarrow Y$ between totally cocomplete $\mathcal{T}$ -categories is cocontinuous if and only if $f^{\text{op}} : X^{\text{op}} \rightarrow Y^{\text{op}}$ is continuous. Hence:

Theorem 7.4. *Taking duals defines an equivalence functor*

$$(-)^{\text{op}} : \mathcal{T}\text{-CoCts} \rightarrow \mathcal{T}\text{-Cts}$$

which commutes with the canonical forgetful functors to $\mathbf{Set}$. Furthermore, $(-)^{\text{op}}$ restricts to separated objects, hence

$$(\mathcal{T}\text{-Cat}_{\text{sep}})^{\mathbb{P}} \simeq (\mathcal{T}\text{-ReprCat}_{\text{sep}})^{\mathbb{V}}.$$

Remark 7.5. We can write the canonical functor $\mathcal{T}\text{-CoCts}_{\text{sep}} \rightarrow \mathbf{Set}^{\mathbb{T}}$ as the composite

$$\mathcal{T}\text{-CoCts}_{\text{sep}} \simeq (\mathcal{T}\text{-ReprCat}_{\text{sep}})^{\mathbb{V}} \rightarrow \mathcal{T}\text{-ReprCat}_{\text{sep}} \rightarrow \mathbf{Set}^{\mathbb{T}},$$

hence its left adjoint sends a $\mathbb{T}$ -algebra X to the totally cocomplete $\mathcal{T}$ -category $\mathbf{V}^X$. Furthermore, we conclude that the monad $\tilde{\mathbf{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on $\mathbf{Set}^{\mathbb{T}}$ (see Remark 4.14) is also induced by the adjunction

$$\mathcal{T}\text{-CoCts}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad} \end{array} \mathbf{Set}^{\mathbb{T}},$$

and that $(\mathcal{T}\text{-ReprCat}_{\text{sep}})^{\mathbb{V}} \simeq (\mathbf{Set}^{\mathbb{T}})^{\tilde{\mathbf{V}}}$.

Examples 7.6. For ordered sets, Theorem 7.4 just states the trivial fact that the category **Sup** of sup-lattices is equivalent to the category **Inf** of inf-lattices. We find it interesting to see that the topological counterpart of this result states that the category **ContLat** of continuous lattices and Scott-continuous and inf-preserving maps is equivalent to the category of Eilenberg–Moore algebras for the upper Vietoris monad on the category **StablyComp** of stably compact spaces and spectral maps. Furthermore, by Remark 7.5, **ContLat** is also equivalent to the category of Eilenberg–Moore algebras for the (classical) Vietoris monad on the category of compact Hausdorff spaces and continuous maps. We note that the latter equivalence was shown in [Wyler, 1981].

For $\mathcal{T} = \mathcal{U}_{\mathbb{P}_+}$, the Eilenberg–Moore category $\mathbf{CompHaus}^{\tilde{\mathbb{V}}}$ of the monad $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on **CompHaus** (see Example 4.15) is equivalent to $\mathbf{Set}^{\mathbb{P}}$, and this category is described in [Gutierrez and Hofmann, 2012] as the category of continuous lattices equipped with an internal action of $[0, \infty]$ and action-preserving morphisms of continuous lattices. A slightly different monad on **CompHaus** one obtains for $\mathcal{T} = \mathcal{U}_{\mathbb{P}_\lambda}$, and the category of Eilenberg–Moore algebras of this monad is equivalent to the category of separated injective objects and left adjoint morphisms in **UApp**.

Remark 7.7. Following [Rosebrugh and Wood, 2004], we consider, for a monad $\mathbb{D}$ on a category $\mathbf{C}$ where idempotents split, the full subcategory $\mathbf{Spl}(\mathbf{C}^{\mathbb{D}})$ of $\mathbf{C}^{\mathbb{D}}$ defined by the *split structures*, that is, by those $\mathbb{D}$ -algebras $(X, \alpha : DX \rightarrow X)$ for which exists a homomorphism $t : X \rightarrow DX$ with $\alpha \cdot t = 1_X$. We put

$$\mathcal{T}\text{-CoCts}_{\text{spl}} := \mathbf{Spl}((\mathcal{T}\text{-Cat})^{\mathbb{P}}) \quad \text{and} \quad \mathcal{T}\text{-Cts}_{\text{spl}} := \mathbf{Spl}((\mathcal{T}\text{-ReprCat})^{\mathbb{V}}).$$

Since our monads are of Kock–Zöberlein type, these splittings are actually adjoint to the algebra structure. Hence, a totally cocomplete separated $\mathcal{T}$ -category X belongs to $\mathcal{T}\text{-CoCts}_{\text{spl}}$ if and only if $\text{Sup}_X : PX \rightarrow X$ has a left adjoint in $\mathcal{T}\text{-Cat}$ (and hence in $\mathcal{T}\text{-CoCts}_{\text{sep}}$), and a totally complete separated $\mathcal{T}$ -category X belongs to $\mathcal{T}\text{-Cts}_{\text{spl}}$ if and only if $\text{Inf}_X : VX \rightarrow X$ has a right adjoint in $\mathcal{T}\text{-ReprCat}$ (and hence in $\mathcal{T}\text{-Cts}_{\text{sep}}$). For X in $\mathcal{T}\text{-CoCts}_{\text{spl}}$, the splitting $t : X \rightarrow PX$ of $\text{Sup}_X : PX \rightarrow X$ is left adjoint and therefore a pseudo-homomorphism; hence, with the help of (5), we see that $V(\alpha^{\text{op}}) \cdot t^{\text{op}}$ is a splitting of $\text{Inf}_{X^{\text{op}}}$ in $(\mathcal{T}\text{-ReprCat})^{\mathbb{V}}$. Therefore the equivalence functor $(-)^{\text{op}} : \mathcal{T}\text{-CoCts} \rightarrow \mathcal{T}\text{-Cts}$ of Theorem 7.4 restricts to a functor

$$(-)^{\text{op}} : \mathcal{T}\text{-CoCts}_{\text{spl}} \rightarrow \mathcal{T}\text{-Cts}_{\text{spl}};$$

however, in general we do not obtain an equivalence as the following example shows.

Example 7.8. We consider the case of topological spaces, that is $\mathcal{T} = \mathcal{U}_2$. For a topological space X , PX is the filter space of X ([Hofmann and Tholen, 2010, Example 4.10]) which is known to be spectral, and so is every split algebra for $\mathbb{P}$. Since the dual of a spectral space is spectral, the image of $(-)^{\text{op}}$ contains only spectral spaces. For a stably compact space X , VX is spectral if and only if X is spectral. In fact, since $h_X : X \rightarrow VX$ is in **StablyComp** and a topological embedding, X is spectral if VX is so. If X is spectral, then the topology of VX is generated by the sets $V^\diamond$ where V runs through all compact opens of X ; and for such V one easily sees that $V^\diamond$ is compact in VX (using Alexander’s Subbase Lemma and $(\bigcup_i V_i)^\diamond = \bigcup_i V_i^\diamond$). Since VX is always a split algebra for $\mathbb{V}$, we conclude that $(-)^{\text{op}}$ is not essentially surjective on objects.

Similarly, for a compact Hausdorff space X , X is a Stone space if and only if $\tilde{V}X$ is a Stone space (if X is Stone, then VX is spectral and hence $\tilde{V}X$ is Stone).

8. THE KLEISLI CATEGORY OF $\mathbb{V}$

Every $\mathcal{T}$ -functor $r : X \rightarrow VY$ gives rise to a $\mathbb{V}$ -matrix $\lrcorner r \rceil : X \dashrightarrow Y$ where $\lrcorner r \rceil(x, y) = r(x)(y)$. In the sequel we are interested in the case where $X = (X, a)$ and $Y = (Y, b)$ are representable $\mathcal{T}$ -categories and $r : X \rightarrow VY$ is a pseudo-homomorphism.

Proposition 8.1. *Let $X = (X, a)$ and $Y = (Y, b)$ be representable $\mathcal{T}$ -categories. Then $r \mapsto \lrcorner r \lrcorner$ defines a bijection between $\mathcal{T}\text{-ReprCat}(X, VY)$ and the subset of $V\text{-Dist}(X_0, Y_0)$ consisting of all those V -distributors $\psi : X_0 \multimap Y_0$ making the diagram*

$$\begin{array}{ccc} T(X_0) & \xrightarrow{T_\xi \psi} & T(Y_0) \\ a \downarrow & & \downarrow b \\ X_0 & \xrightarrow[\psi]{} & Y_0 \end{array} \quad (6)$$

commutative.

Proof. Let $\alpha : TX \rightarrow X$ and $\beta : TY \rightarrow Y$ be pseudo-algebra structures of X and Y respectively. Assume first that $r : X \rightarrow VY$ is a homomorphism. Then $\lrcorner r \lrcorner$ is a $\mathcal{T}$ -functor $\lrcorner r \lrcorner : X^{\text{op}} \otimes Y \rightarrow V$ and hence also a V -functor $\lrcorner r \lrcorner : X_0^{\text{op}} \otimes Y_0 \rightarrow V$. But the latter is equivalent to $\lrcorner r \lrcorner$ being a V -distributor $\lrcorner r \lrcorner : X_0 \multimap Y_0$. Furthermore, for all $\mathfrak{x} \in TX$ and $y \in Y$,

$$\mu_Y \cdot Tr(\mathfrak{x})(y) = b \cdot T_\xi \text{ev}_Y(Tr(\mathfrak{x}), y) = b \cdot T_\xi \lrcorner r \lrcorner(\mathfrak{x}, y);$$

hence $r \cdot \alpha = \mu_Y \cdot Tr$ if and only if $\lrcorner r \lrcorner \cdot a = b \cdot T_\xi \lrcorner r \lrcorner$ (note that $\lrcorner r \lrcorner \cdot \alpha = \lrcorner r \lrcorner \cdot a_0 \cdot \alpha = \lrcorner r \lrcorner \cdot a$ since $\lrcorner r \lrcorner$ is a V -distributor).

Assume now that $\psi : X \multimap Y$ is a V -distributor making the diagram (6) commutative. Then, for every $x \in X$,

$$b \cdot T_\xi(\psi \cdot x) \cdot e_1 = \psi \cdot a \cdot e_X \cdot x = \psi \cdot x,$$

hence $\psi \cdot x$ can be seen a $\mathcal{T}$ -distributor of type $G \multimap Y$. We conclude that $\psi = \lrcorner r \lrcorner$ for $r : X \rightarrow VY$, $x \mapsto \psi \cdot x$. Finally, r is a V -functor since ψ is a V -distributor, and r is a homomorphism by the considerations at the end of the first part of the proof. $\square$

Remark 8.2. For a V -distributor $\psi : X_0 \multimap Y_0$, commutativity of (6) is equivalent to $\psi \cdot \alpha = b \cdot T_\xi \psi$ since $\psi \cdot a_0 = \psi$. Hence, if Y is of the form $Y = (Y, \beta : TY \rightarrow Y)$, (6) commutes if and only if $\psi \cdot \alpha = \beta \cdot T_\xi \psi$. Furthermore, for every pseudo-homomorphism $f : X \rightarrow Y$, the V -distributor $f_* : X_0 \multimap Y_0$ is a morphism $f_* : X \multimap Y$ in $\mathcal{T}\text{-ReprDist}$ since

$$f_* \cdot \alpha_* = (f \cdot \alpha)_* = (\beta \cdot Tf)_* = b \cdot T(f_*).$$

More generally, given a V -functor $f : X_0 \rightarrow Y_0$, the V -distributor $f_* : X_0 \multimap Y_0$ makes (6) commutative if and only if $f : X \rightarrow Y$ is a pseudo-homomorphism.

We write $\mathcal{T}\text{-ReprDist}$ for the category with objects all representable $\mathcal{T}$ -categories, and a morphisms $\psi : X \multimap Y$ in $\mathcal{T}\text{-ReprDist}$ is a V -distributor $\psi : X_0 \multimap Y_0$ making (6) commutative. Composition in $\mathcal{T}\text{-ReprDist}$ is given by V -relational composition, and $a_0 : X \multimap X$ is the identify arrow on X . Hence, $(X, a) \mapsto (X, a_0)$ defines a faithful functor

$$\mathcal{T}\text{-ReprDist} \rightarrow V\text{-Dist}.$$

The following lemma is obvious.

Lemma 8.3. *Let $X = (X, a)$, $Y = (Y, b)$ and $Z = (Z, c)$ be in $\mathcal{T}\text{-ReprDist}$ and let $\varphi : (X, a_0) \rightarrow (Y, b_0)$ and $\psi : (Y, b_0) \rightarrow (Z, c_0)$ be V -distributors. If $\psi \cdot \varphi : X \multimap Z$ and $\psi : Y \multimap Z$ are actually morphisms in $\mathcal{T}\text{-ReprDist}$ and ψ is mono in $V\text{-Dist}$, then $\varphi : X \multimap Y$ is in $\mathcal{T}\text{-ReprDist}$.*

By definition, $f : X \rightarrow Y$ in $\mathcal{T}\text{-ReprCat}$ is downwards open (see Definition 4.16) precisely if $a \cdot T_\xi(f^*) = f^* \cdot b$, and therefore:

Proposition 8.4. *The following assertions are equivalent, for $f : X \rightarrow Y$ in $\mathcal{T}\text{-ReprCat}$.*

- (i) *f is downwards open.*
- (ii) *The V -distributor $f^* : Y_0 \multimap X_0$ makes (6) commutative (that is, $f^* : Y \multimap X$ is a morphism in $\mathcal{T}\text{-ReprDist}$).*
- (iii) *$f_* : X \multimap Y$ is left adjoint in $\mathcal{T}\text{-ReprDist}$.*

Theorem 8.5. *The Kleisli category $\mathcal{T}\text{-ReprCat}_{\mathbb{V}}$ of $\mathbb{V}$ is equivalent to $\mathcal{T}\text{-ReprDist}$.*

Proof. It is left to show that $r \mapsto \lrcorner r \lrcorner$ preserves composition. To see this, let $r : X \rightarrow VY$ and $s : Y \rightarrow VZ$ be in $\mathcal{T}\text{-ReprCat}_{\mathbb{V}}$. First note that, for every $y \in Y$ and $z \in Z$,

$$h_Z^\circ \cdot s_*(y, z) = s_*(y, h_Z(z)) = [h_Z(z), s(y)] = s(y)(z) = \lrcorner s \lrcorner(y, z),$$

and therefore $\omega_Y \cdot Vs : VY \rightarrow VZ$ sends $\varphi : G \multimap Y$ to $\lrcorner s \lrcorner \cdot \varphi$ (see Remark 4.9). Consequently,

$$\lrcorner s \lrcorner \cdot \lrcorner r \lrcorner(x, z) = \lrcorner s \lrcorner \cdot \lrcorner r \lrcorner \cdot x(z) = \omega_Y \cdot Vs \cdot r(x)(z),$$

for all $x \in X$ and $z \in Z$. □

Corollary 8.6. *The functor $(-)_* : \mathcal{T}\text{-ReprCat} \rightarrow \mathcal{T}\text{-ReprDist}$ is left adjoint to*

$$\mathcal{T}\text{-ReprDist} \rightarrow \mathcal{T}\text{-ReprCat}, (\psi : X \multimap Y) \mapsto (\psi \cdot - : VX \rightarrow VY).$$

Here we think of an element $\varphi \in VX$ as a morphism $\varphi : G \multimap X$ in $\mathcal{T}\text{-ReprDist}$. The units and counits are given by $h_X : X \rightarrow VX$ and $h_X^* : VX \multimap X$ respectively.

Remark 8.7. Certainly, the adjunction above can be restricted to separated $\mathcal{T}$ -categories to yield

$$\mathcal{T}\text{-ReprDist}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad (-)_* \quad} \end{array} \mathcal{T}\text{-ReprCat}_{\text{sep}}.$$

Furthermore, the monad $\tilde{\mathbb{V}} = (\tilde{V}, \tilde{\omega}, \tilde{h})$ on $\mathbf{Set}^{\mathbb{T}}$ of Remark 4.14 is also induced by the composite adjunction

$$\mathcal{T}\text{-ReprDist}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad \quad} \end{array} \mathcal{T}\text{-ReprCat}_{\text{sep}} \begin{array}{c} \xrightarrow{\quad \top \quad} \\ \xleftarrow{\quad \quad} \end{array} \mathbf{Set}^{\mathbb{T}}.$$

The fully faithful comparison functor $(\mathbf{Set}^{\mathbb{T}})_{\tilde{\mathbb{V}}} \rightarrow \mathcal{T}\text{-ReprDist}_{\text{sep}}$ induces an equivalence between the Kleisli category $(\mathbf{Set}^{\mathbb{T}})_{\tilde{\mathbb{V}}}$ of $\tilde{\mathbb{V}}$ and the full subcategory $\mathcal{T}\text{-ReprDist}_{\text{sep}}$ of $\mathcal{T}\text{-ReprDist}$ defined by objects of the form $X = (X, \alpha : TX \rightarrow X)$ (i.e. where X_0 is a discrete $\mathbb{V}$ -category).

Example 8.8. An *Esakia space* [Esakia, 1974] is a Priestley space $(X, \leq, \alpha)$ where the down-closure of every open (with respect to α) subset $A \subseteq X$ is again open (with respect to α or, equivalently, with respect to $a \leq \cdot \alpha$). We find it worthwhile to mention that this condition just states that $i : (X, \alpha) \rightarrow (X, \leq \cdot \alpha)$, $x \mapsto x$ is downwards open. A *morphism of Esakia spaces* (also called bounded morphism or p-morphism) is a homomorphism $f : (X, \leq, \alpha) \rightarrow (Y, \leq, \beta)$ such that, for all $x \in X$ and $y \in Y$ with $f(x) \leq y$, there is some $x' \in X$ with $x \leq x'$ and $f(x') = y$; and this condition just means that the diagram

$$\begin{array}{ccc} (X, \leq \cdot \alpha) & \xrightarrow{i_X^*} & (X, \alpha) \\ f_* \downarrow \circ & & \downarrow \circ f \\ (Y, \leq \cdot \beta) & \xrightarrow{i_Y^*} & (Y, \beta) \end{array}$$

commutes.

Motivated by the example above, we introduce the following notion.

Definition 8.9. A separated representable $\mathcal{T}$ -category (X, a) (with algebra structure $\alpha : TX \rightarrow X$) is an *Esakia $\mathcal{T}$ -category* whenever $i : (X, \alpha) \rightarrow (X, a)$ is downwards open.

Proposition 8.10. *The following assertions are equivalent, for a separated representable $\mathcal{T}$ -category $X = (X, a)$.*

- (i) *X is an Esakia $\mathcal{T}$ -category.*

- (ii) The $\mathbf{V}$ -relation $a_0 : X \rightarrowtail X$ is a morphism $a_0 : (X, a) \multimap (X, \alpha)$ in $\mathcal{T}\text{-ReprDist}$, that is, the diagram

$$\begin{array}{ccc} TX & \xrightarrow{T_\xi a_0} & TX \\ \alpha \downarrow & & \downarrow \alpha \\ X & \xrightarrow{a_0} & X \end{array} \quad (7)$$

commutes in $\mathbf{V}\text{-Dist}$.

- (iii) X is a split subobject in $\mathcal{T}\text{-ReprDist}$ of a $\mathbb{T}$ -algebra $(Y, \beta) \in \mathbf{Set}^{\mathbb{T}}$.

Proof. The equivalence (i) $\Leftrightarrow$ (ii) is clear by definition since $i^* = a_0$, and (ii) $\Rightarrow$ (iii) follows from $i_* \cdot i^* = a_0$. To see (iii) $\Rightarrow$ (ii), let (Y, β) be a $\mathbb{T}$ -algebra and $\psi : (X, a) \multimap (Y, \beta)$, $\varphi : (Y, \beta) \multimap (X, a)$ be in $\mathcal{T}\text{-ReprDist}$ with $\varphi \cdot \psi = a_0$. Then

$$\psi \cdot \varphi = (\psi \cdot i_*) \cdot (i^* \cdot \varphi)$$

in $\mathbf{V}\text{-Dist}$, $\psi \cdot \varphi$, $\psi \cdot i_*$ are in $\mathcal{T}\text{-ReprDist}$ and $\psi \cdot i_*$ is mono in $\mathbf{V}\text{-Dist}$, hence, by Lemma 8.3, $i^* \cdot \varphi$ is in $\mathcal{T}\text{-ReprDist}$. Consequently, $i^* = i^* \cdot \varphi \cdot \psi$ is in $\mathcal{T}\text{-ReprDist}$. $\square$

Example 8.11. We return to the case of topological spaces. By Example 7.8, the monad $\mathbb{W}$ on $\mathbf{StablyComp}$ restricts to $\mathbf{Spec}$, and the Kleisli category $\mathbf{Spec}_{\mathbb{W}}$ corresponds to the full subcategory $\mathbf{SpecDist}$ of $\mathcal{U}_2\text{-ReprDist}$ defined by all spectral spaces. Similarly, the monad $\tilde{\mathbb{W}}$ on $\mathbf{CompHaus}$ restricts to $\mathbf{Stone}$ and the Kleisli category $\mathbf{Stone}_{\tilde{\mathbb{W}}}$ corresponds to the full subcategory $\mathbf{StoneDist}$ of $\mathbf{SpecDist}$ defined by all Stone spaces. Then $\mathbf{StoneDist}$ is dually equivalent to the category $\mathbf{Bool}_{\perp, \vee}$ of Boolean algebras and finite suprema preserving maps (see [Halmos, 1962; Sambin and Vaccaro, 1988]), and $\mathbf{SpecDist}$ is dually equivalent to the category $\mathbf{DLat}_{\perp, \vee}$ of distributive lattices and finite suprema preserving maps (see [Cignoli *et al.*, 1991]). Furthermore, these dualities are closely related to duality results for Boolean algebras with operator (see [Kupke *et al.*, 2004]) and distributive lattices with operator (see [Petrovich, 1996; Bonsangue *et al.*, 2007]). One easily verifies that $\mathbf{DLat}_{\perp, \vee}$ is idempotent split complete (since it is a full subcategory of the algebraic category of sup-semilattices and homomorphisms and it is closed there under split quotients), and therefore also $\mathbf{SpecDist}$ is so. Consequently, by Proposition 8.10, the full subcategory of $\mathbf{SpecDist}$ defined by all Esakia spaces is the idempotent split completion of $\mathbf{StoneDist}$; which, by Esakia duality [Esakia, 1974], then implies that the category $\mathbf{Heyt}_{\perp, \vee}$ of Heyting algebras and finite suprema preserving maps is the idempotent split completion of $\mathbf{Bool}_{\perp, \vee}$.

A morphism $f : X \rightarrow Y$ of Esakia $\mathcal{T}$ -categories $X = (X, a)$ and $Y = (Y, b)$ (where $b = b_0 \cdot \beta$) is a homomorphism making

$$\begin{array}{ccc} (X, a) & \xrightarrow{i_X^*} & (X, \alpha) \\ f_* \downarrow \circ & & \downarrow \circ f \\ (Y, b) & \xrightarrow{i_Y^*} & (Y, \beta) \end{array}$$

commutative in $\mathcal{T}\text{-ReprDist}$, which can be equivalently expressed by saying that either of the diagrams

$$\begin{array}{ccc} X & \xrightarrow{\ulcorner a_0 \urcorner} & V(X, \alpha) \\ f \downarrow & & \downarrow Vf \\ Y & \xrightarrow{\ulcorner b_0 \urcorner} & V(Y, \beta) \end{array} \quad \text{or} \quad \begin{array}{ccc} (X, \alpha) & \xrightarrow{\ulcorner a_0 \urcorner} & \tilde{V}(X, \alpha) \\ f \downarrow & & \downarrow \tilde{V}f \\ (Y, \beta) & \xrightarrow{\ulcorner b_0 \urcorner} & \tilde{V}(Y, \beta) \end{array}$$

commutes in $\mathcal{T}\text{-ReprCat}$ or $\mathbf{Set}^{\mathbb{T}}$ respectively. Let now $X = (X, \alpha)$ be in $\mathbf{Set}^{\mathbb{T}}$ and $r : X \rightarrow \tilde{V}X$ be a homomorphism. Then, with $a_0 := \ulcorner r \urcorner : X \rightarrowtail X$, the diagram (7) commutes by Proposition 8.1. Therefore $(X, a_0 \cdot \alpha)$ is an Esakia $\mathcal{T}$ -category provided that a_0 is a separated $\mathbf{V}$ -category

structure on the set X . Summing up, we have identified the category of Esakia $\mathcal{T}$ -categories and morphisms as the full subcategory of the category $\mathbf{Coalg}(\tilde{V})$ of coalgebras for $\tilde{V} : \mathbf{Set}^{\mathbb{T}} \rightarrow \mathbf{Set}^{\mathbb{T}}$ defined by those coalgebras $r : X \rightarrow \tilde{V}X$ whose mate $a_0 := \lrcorner r \lrcorner : X \multimap X$ is a separated V -category structure on the set X . This observation represents a generalisation of [Davey and Galati, 2003].

REFERENCES

- BANASCHEWSKI, B., LOWEN, R. and VAN OLMEN, C. (2006), Sober approach spaces, *Topology Appl.* **153** (16), 3059–3070.
- BARR, M. (1970), Relational algebras, in *Reports of the Midwest Category Seminar, IV*, pages 39–55, Lecture Notes in Mathematics, Vol. 137. Springer, Berlin.
- BETTI, R., CARBONI, A., STREET, R. and WALTERS, R. (1983), Variation through enrichment, *J. Pure Appl. Algebra* **29** (2), 109–127.
- BONSANGUE, M. M., KURZ, A. and REWITZKY, I. M. (2007), Coalgebraic representations of distributive lattices with operators, *Topology Appl.* **154** (4), 778–791.
- CIGNOLI, R., LAFALCE, S. and PETROVICH, A. (1991), Remarks on Priestley duality for distributive lattices, *Order* **8** (3), 299–315.
- CLEMENTINO, M. M. and HOFMANN, D. (2003), Topological features of lax algebras, *Appl. Categ. Structures* **11** (3), 267–286.
- CLEMENTINO, M. M. and HOFMANN, D. (2004), Effective descent morphisms in categories of lax algebras, *Appl. Categ. Structures* **12** (5-6), 413–425.
- CLEMENTINO, M. M. and HOFMANN, D. (2009a), Lawvere completeness in topology, *Appl. Categ. Structures* **17** (2), 175–210, [arXiv:math.CT/0704.3976](#).
- CLEMENTINO, M. M. and HOFMANN, D. (2009b), Relative injectivity as cocompleteness for a class of distributors, *Theory Appl. Categ.* **21**, No. 12, 210–230, [arXiv:math.CT/0807.4123](#).
- CLEMENTINO, M. M., HOFMANN, D. and THOLEN, W. (2003), Exponentiability in categories of lax algebras, *Theory Appl. Categ.* **11**, No. 15, 337–352.
- CLEMENTINO, M. M. and THOLEN, W. (1997), A characterization of the Vietoris topology, in *Proceedings of the 12th Summer Conference on General Topology and its Applications (North Bay, ON, 1997)*, volume 22, pages 71–95.
- CLEMENTINO, M. M. and THOLEN, W. (2003), Metric, topology and multicategory—a common approach, *J. Pure Appl. Algebra* **179** (1-2), 13–47.
- DAVEY, B. A. and GALATI, J. C. (2003), A coalgebraic view of Heyting duality, *Studia Logica* **75** (3), 259–270.
- DAY, B. J. and KELLY, G. M. (1970), On topologically quotient maps preserved by pullbacks or products, *Proc. Cambridge Philos. Soc.* **67**, 553–558.
- EILENBERG, S. and KELLY, G. M. (1966), Closed categories, in *Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965)*, pages 421–562, Springer, New York.
- ESAKIA, L. (1974), Topological Kripke models, *Dokl. Akad. Nauk SSSR* **214**, 298–301.
- ESCARDÓ, M. H. (1998), Properly injective spaces and function spaces, *Topology Appl.* **89** (1-2), 75–120.
- GIERZ, G., HOFMANN, K. H., KEIMEL, K., LAWSON, J. D., MISLOVE, M. W. and SCOTT, D. S. (1980), *A compendium of continuous lattices*, Springer-Verlag, Berlin, xx+371 pages.
- GUTIERRES, G. and HOFMANN, D. (2012), Approaching metric domains, *accepted for publication in Appl. Categ. Structures* [arXiv:math.GN/1103.4744](#).
- HALMOS, P. R. (1962), *Algebraic logic*, Chelsea Publishing Co., New York, 271 pages.
- HERMIDA, C. (2000), Representable multicategories, *Adv. Math.* **151** (2), 164–225.
- HOCHSTER, M. (1969), Prime ideal structure in commutative rings, *Trans. Amer. Math. Soc.* **142**, 43–60.

- HOFMANN, D. (2007), Topological theories and closed objects, *Adv. Math.* **215** (2), 789–824.
- HOFMANN, D. (2010), Duality for distributive spaces, Technical report, [arXiv:math.CT/1009.3892](#).
- HOFMANN, D. (2011), Injective spaces via adjunction, *J. Pure Appl. Algebra* **215** (3), 283–302, [arXiv:math.CT/0804.0326](#).
- HOFMANN, D. and STUBBE, I. (2011), Towards Stone duality for topological theories, *Topology Appl.*, **158** (7), 913–925, [arXiv:math.CT/1004.0160](#).
- HOFMANN, D. and THOLEN, W. (2010), Lawvere completion and separation via closure, *Appl. Categ. Structures* **18** (3), 259–287, [arXiv:math.CT/0801.0199](#).
- HOFMANN, D. and WASZKIEWICZ, P. (2011), Approximation in quantale-enriched categories, *Topology Appl.* **158** (8), 963–977, [arXiv:math.CT/1004.2228](#).
- ISELL, J. R. (1975), Meet-continuous lattices, in *Symposia Mathematica, Vol. XVI (Convegno sui Gruppi Topologici e Gruppi di Lie, INDAM, Rome, 1974)*, pages 41–54, Academic Press, London.
- ISELL, J. R. (1986), General function spaces, products and continuous lattices, *Math. Proc. Cambridge Philos. Soc.* **100** (2), 193–205.
- JOHNSTONE, P. T. (1986), *Stone spaces*, volume 3 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, xxii+370 pages, reprint of the 1982 edition.
- JUNG, A. (2004), Stably compact spaces and the probabilistic powerspace construction, in J. Desharnais and P. Panangaden, editors, *Domain-theoretic Methods in Probabilistic Processes*, volume 87, 15pp.
- KELLY, G. M. (1982), *Basic concepts of enriched category theory*, volume 64 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge, 245 pages, also in: *Repr. Theory Appl. Categ.* No. 10 (2005), 1–136.
- KUPKE, C., KURZ, A. and VENEMA, Y. (2004), Stone coalgebras, *Theoret. Comput. Sci.* **327** (1-2), 109–134.
- LAWSON, J. (2011), Stably compact spaces, *Math. Structures Comput. Sci.* **21** (1), 125–169.
- LAWVERE, F. W. (1973), Metric spaces, generalized logic, and closed categories, *Rendiconti del Seminario Matematico e Fisico di Milano* **43**, 135–166, Republished in: *Reprints in Theory and Applications of Categories*, No. 1 (2002) pp 1–37.
- LINTON, F. E. J. (1969), Coequalizers in categories of algebras, in *Sem. on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67)*, pages 75–90, Springer, Berlin.
- LOWEN, R. (1989), Approach spaces: a common supercategory of TOP and MET, *Math. Nachr.* **141**, 183–226.
- LOWEN, R. (1997), *Approach spaces*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, x+253 pages, the missing link in the topology-uniformity-metric triad, Oxford Science Publications.
- LUCYSHYN-WRIGHT, R. B. B. (2011), Domains occur among spaces as strict algebras among lax, *Math. Structures Comput. Sci.* **21** (3), 647–670.
- MANES, E. G. (1969), A triple theoretic construction of compact algebras., *Sem. on Triples and Categorical Homology Theory, ETH Zürich 1966/67, Lect. Notes Math.* 80, 91–118 (1969).
- MANES, E. G. (1976), *Algebraic theories*, Springer-Verlag, New York, vii+356 pages, graduate Texts in Mathematics, No. 26.
- NACHBIN, L. (1950), *Topologia e Ordem*, Univ. of Chicago Press, in Portuguese, English translation: *Topology and Order*, Van Nostrand, Princeton (1965).
- PETROVICH, A. (1996), Distributive lattices with an operator, *Studia Logica* **56** (1-2), 205–224, special issue on Priestley duality.
- PISANI, C. (1999), Convergence in exponentiable spaces, *Theory Appl. Categ.* **5**, No. 6, 148–162.
- PRIESTLEY, H. A. (1970), Representation of distributive lattices by means of ordered stone spaces, *Bull. London Math. Soc.* **2**, 186–190.

- ROSEBRUGH, R. and WOOD, R. J. (1994), Constructive complete distributivity. IV, *Appl. Categ. Structures* **2** (2), 119–144.
- ROSEBRUGH, R. and WOOD, R. J. (2004), Split structures, *Theory Appl. Categ.* **13**, No. 12, 172–183.
- SAMBIN, G. and VACCARO, V. (1988), Topology and duality in modal logic, *Ann. Pure Appl. Logic* **37** (3), 249–296.
- SCHUBERT, C. and SEAL, G. J. (2008), Extensions in the theory of lax algebras, *Theory Appl. Categ.* **21**, No. 7, 118–151.
- SCHWARZ, F. (1984), Product compatible reflectors and exponentiability, in *Categorical topology (Toledo, Ohio, 1983)*, volume 5 of *Sigma Ser. Pure Math.*, pages 505–522, Heldermann, Berlin.
- SEAL, G. J. (2005), Canonical and op-canonical lax algebras, *Theory Appl. Categ.* **14**, 221–243.
- SEAL, G. J. (2009), A Kleisli-based approach to lax algebras, *Appl. Categ. Structures* **17** (1), 75–89.
- SIMMONS, H. (1982), A couple of triples, *Topology Appl.* **13** (2), 201–223.
- STONE, M. H. (1936), The theory of representations for Boolean algebras, *Trans. Amer. Math. Soc.* **40** (1), 37–111.
- STONE, M. H. (1938), Topological representations of distributive lattices and Brouwerian logics, *Časopis pro pěstování matematiky a fysiky* **67** (1), 1–25, <http://dml.cz/handle/10338.dmlcz/124080>.
- THOLEN, W. (2009), Ordered topological structures, *Topology Appl.* **156** (12), 2148–2157.
- VAN OLMEN, C. (2005), *A study of the interaction between frame theory and approach theory*, Ph.D. thesis, University of Antwerp.
- VIETORIS, L. (1922), Bereiche zweiter Ordnung, *Monatsh. Math. Phys.* **32** (1), 258–280.
- WOOD, R. (2004), Ordered sets via adjunction, in *Categorical foundations: special topics in order, topology, algebra, and sheaf theory*, volume 97 of *Encyclopedia Math. Appl.*, pages 5–47, Cambridge Univ. Press, Cambridge.
- WYLER, O. (1981), Algebraic theories of continuous lattices, Continuous lattices, Proc. Conf., Bremen 1979, Lect. Notes Math. 871, 390–413 (1981).

CIDMA – CENTER FOR RESEARCH AND DEVELOPMENT IN MATHEMATICS AND APPLICATIONS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, 3810-193 AVEIRO, PORTUGAL

E-mail address: `dirk@ua.pt`